


Product List

by DaeHan New Pharm Co., Ltd.



Cardiovascular agents

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Cilotal Tab.		Cilostazol 50mg	Reduction of the symptoms of intermittent claudication	100mg orally twice a day administered at least 30 minutes before or 2 hours after breakfast and dinner	30T 500T
Newpido Tab.		Clopidogrel 75mg	Arterial thromboembolism prophylaxis (myocardial infarction prophylaxis, stroke prophylaxis, thrombosis prophylaxis)	- Acute Coronary Syndrome : 300mg oral loading dose and then continue at 75mg once daily - Recent MI, Recent Stroke, or Established Peripheral Arterial Disease : 75mg once daily orally without a loading dose	30T 100T
Metazin Tab.		Trimetazidine HCl 20mg	Angina pectoris	1 tablet, 3 times daily	30T 300T
Lapine Tab.		Lacidipine 4mg	Hypertension	starting dose : 2 mg, once daily (Daily maximum dose is 6mg)	90T
DHNP Losartan plus Tab.		Losartan potassium 50mg Hydrochlorothiazide 12.5mg	Hypertension	1 tablet, once daily	30T 300T
NP Forge Tab. 5/80mg		Valsartan 80mg Amlodipine 5mg	Hypertension	1 tablet, once daily	28T
NP Forge Tab. 5/160mg		Valsartan 160mg Amlodipine 5mg			
NP Forge Tab. 10/160mg		Valsartan 160mg Amlodipine 10mg			
NP Star Tab. 40/5mg		Telmisartan 40mg Amlodipine 5mg	Hypertension	1 tablet, once daily regardless of meals	30T
NP Star Tab. 40/10mg		Telmisartan 40mg Amlodipine 10mg			
NP Star Tab. 80/5mg		Telmisartan 80mg Amlodipine 5mg			
Lovatadin Tab.		Lovastatin 20mg	Primary hypercholesterolemia : specifically hyperlipoproteinemia types IIa or IIb	10 to 80mg/day in single or 2 divided doses (Daily maximum dose is 80mg)	30T 500T
Newtor Tab. 10mg		Atorvastatin calcium 10.84mg (As Atorvastatin 10mg)	1. Hypercholesterolemia, hyperlipoproteinemia, and/or hypertriglyceridemia 2. Reduction of reducing the risk of cardiovascular events (myocardial infarction prophylaxis, stroke prophylaxis)	- Initial dose : 10, 20 or 40mg, once daily - Maintenance dose : 10 to 80mg, once daily	30T 100T
Newtor Tab. 20mg		Atorvastatin calcium 21.69mg (As Atorvastatin 20mg)			
Newsto Tab. 10mg		Rosuvastatin calcium 10.40mg (As Rosuvastatin 10mg)	Hypercholesterolemia, hyperlipoproteinemia, and/or hypertriglyceridemia	- Initial dose : 5mg, once daily - Maintenance dose : 10mg, once daily	28T

Cardiovascular agents

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Enpitorin Tab. 10/10mg		Simvastatin 10mg Ezetimibe 10mg	1. Primary (heterozygous familial and nonfamilial) hypercholesterolemia 2. Mixed hyperlipoproteinemia, homozygous familial hypercholesterolemia (HoFH)	1 tablet, once daily in the evening	30T
Enpitorin Tab. 10/20mg		Simvastatin 20mg Ezetimibe 10mg			

Endocrine system agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Grizide Tab.		Gliclazide 80mg	Insulin non-dependent diabetes mellitus	Initial dose : 40~80mg, 1-2 times daily	30T 500T
Gerimet Tab.		Glimepiride 2mg Metformin HCl 500mg	Insulin non-dependent diabetes mellitus	1 tablet, 1-2 times daily	30T 100T

Neuropsychiatric agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Enpicept Tab. 10mg		Donepezil HCl 10mg	Alzheimer's disease	5~10mg, once daily before bedtime	30T
Enpicept Tab. 5mg		Donepezil HCl 5mg			
Sporatin Tab.		Acetyl L-carnitine HCl 590mg (As Acetyl L-carnitine 500mg)	1. Primary degenerative cerebral disease 2. Degenerative disease caused by cerebrovascular disease	1 tablet, 2-3 times daily	60T

Hormones

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Hytisone Tab.		Hydrocortisone 10mg	1. Adrenocortical insufficiency : adrenal hyperplasia, primary adrenocortical insufficiency (Addison's Disease), or secondary adrenocortical insufficiency 2. Rheumatic disorders : acute gouty arthritis, ankylosing spondylitis, rheumatoid arthritis, post-traumatic osteoarthritis, pseudogout, or psoriatic arthritis	10~120mg, once daily in 1-4 divided doses	30T 500T

Gastrointestinal agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
S-oem Tab, 20mg		Esomeprazole 20mg	1. Gastroesophageal reflux disease (GERD), 2. Helicobacter pylori (H. pylori) eradication in combination with antimicrobial agent 3. Short-term treatment of frequent dyspepsia or pyrosis (heartburn) 4. NSAID-induced ulcer prophylaxis or healing (gastric) 5. Zollinger–Ellison syndrome 6. Upper GI rebleeding prophylaxis after therapeutic endoscopy	- 20~40mg, once daily - Zollinger–Ellison syndrome : 40~80mg, twice daily	28T
S-oem Tab, 40mg		Esomeprazole 40mg			28T
DHNP Omprazole Cap.		Omeprazole 20mg	1.Short-term treatment of frequent dyspepsia or pyrosis (heartburn) 2. Gastroesophageal reflux disease (GERD) 3. Duodenal ulcer 4. Long-term treatment of gastric hypersecretory conditions, including Zollinger–Ellison syndrome, systemic mastocytosis, and multiple endocrine adenoma syndrome 5. NSAID-induced ulcer prophylaxis	1 capsule, once daily	28C
Rapizole Tab, 10mg		Rabeprazole sodium 10mg	1. Duodenal ulcer 2. Gastroesophageal reflux disease (GERD) 3. Pathological hypersecretory conditions including Zollinger–Ellison syndrome 4. Helicobacter pylori (H. pylori) eradication in the treatment of patients with duodenal ulcer disease, gastric ulcer, dyspepsia, or gastric mucosa associated lymphoid tissue (MALT) lymphoma	- Duodenal ulcer : 10mg, once daily - GERD : 10mg~20mg, once daily - Zollinger–Ellison syndrome : 60mg, once daily - Helicobacter pylori (H. pylori) eradication : 20mg, twice daily with amoxicillin (1,000 mg twice daily) and clarithromycin (500 mg twice daily) for 7 days	28T
Rapizole Tab, 20mg		Rabeprazole sodium 20mg			28T
Camillan Tab.		Ranitidine HCl 168mg (As Ranitidine 150mg)	1. Peptic ulcer disease (duodenal ulcer or gastric ulcer) or gastritis 2. Gastroesophageal reflux disease (GERD) 3. Pathologic GI hypersecretory conditions such as Zollinger–Ellison syndrome, systemic mastocytosis, or multiple endocrine adenoma syndrome 4. Erosive esophagitis	1 tablet, once daily	30T 500T
Newzantin Cap, 150mg		Nizatidine 150mg	1. Gastroesophageal reflux disease (GERD), including endoscopically diagnosed esophagitis (erosive and ulcerative esophagitis) and GERD-associated pyrosis (heartburn). 2. Duodenal ulcer 3. Active benign gastric ulcer	300mg, once daily or 150mg, twice daily (morning and night)	30C 300C
JS Cimetidine Tab.		Cimetidine 200mg	1. Peptic ulcer (duodenal ulcer or gastric ulcer) 2. Gastroesophageal reflux disease (GERD) or esophagitis associated with gastroesophageal reflux disease (GERD) 3. NSAID-induced ulcer prophylaxis 4. Pathological hypersecretory conditions including Zollinger–Ellison syndrome, systemic mastocytosis, and multiple endocrine adenoma syndrome	2 tablets, twice daily or 4 tablets, once daily at bedtime	1000T
Mucoran Tab.		Rebamipide 100mg	1. Gastric ulcer 2. Improvement of gastric mucosal lesions (erosion, bleeding, flare edema) in the following disease : acute gastritis, acute exacerbation of chronic gastritis	1 tablet, 3 times daily (Gastric ulcer: morning, night, before bedtime)	30T 500T
Levoran Tab.		Levosulpiride 25mg	Functional dyspepsia : indigestion, abdominal fullness, vomiting and irritable bowel syndrome	1 tablet, 3 times daily	30T 500T
Mosagatin Tab.		Mosapride citrate 5mg	Gastrointestinal symptoms associated with functional dyspepsia : acid reflux, irritable bowel syndrome and indigestion	1 tablets, 3 times daily (before or after meals)	30T 300T
NP Ren Tab.		Artemisiae argyi folium 95% ethanol ext.(20→1)	1. Gastritis 2. NSAID-induced Gastritis prophylaxis	1 tablet, 3 times daily	30T 500T

Gastrointestinal agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Acitoba Cap. 300mg		Lactobacillus acidophilus 300mg (1.0 X 10 ⁸ as viable cell number s)	Acute diarrhea, irritable bowel syndrome, diarrhea by function colic disease	- Adults and children over 12 years : 300mg, 1~3 times daily - Children 8~12 years : 300mg, 1~2 times daily	30C 500C
Newcon Tab.		Calcium polycarbophil 625mg	Constipation, irritable bowel syndrome, diarrhea	2 tablets, 1~4 times daily	30T 500T
Tambutine Tab, 200mg		Trimebutine maleate 200mg	Digestive function disorder in gastroesophageal reflux, hiatal hernia, gastroduodenitis and gastroduodenitis and gastroduodenal ulcer	1/2~1 capsule, 3 times daily	30T 500T

Musculoskeletal agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Epison Tab.		Eperisone HCl 50mg	1. Painful muscle spasm : scapula symptom, periarthritis, back pain. 2. Spastic paralysis associated with nervous system disease	1 tablet, 3 times daily	30T 500T

Osteoporosis agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Cerobone Plus Tab.		Cholecalciferol concentrated powder 56mg (As Cholecalciferol 5,600 I.U) Risedronate sodium 35mg	1. Female post menopause osteoporosis (treatment and prevention) 2. Male osteoporosis	1 tablet, once a week	4T
Alobone Plus D Tab.		Cholecalciferol concetrate powder 53.33mg (As Vitamin D ₃ 5,600 I.U) Alendronate sodium 91.37mg (As Alendronic acid 70mg)	1. Female post menopause osteoporosis 2. Male osteoporosis	1 tablet, once a week	4T
Enpibone Inj.		Ibandronate sodium monohydrate 3.375mg/3ml (As Ibandronic acid 1mg/mL)	Female post menopause osteoporosis	Intravenous injection for 15~30 seconds of 3mg every 3 months	1syr.

Respiratory system agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Esten Cap. 300mg		Erdosteine 300mg	Mucolytic and expectorant agent for acute and chronic respiratory disease	1 capsule, 2~3 times daily	30C 500C
Elpizin Tab.		Levodropropizine 60mg	Acute and chronic bronchitis	1 tablet, 3 times daily	30T 300T
Mucidin Cap. 200mg		Acetylcysteine 200mg	Mucolysis of viscous or inspissated mucous secretions in patients with pulmonary conditions (pneumonia, bronchitis, cystic fibrosis, atelectasis due to mucous obstruction, and post-traumatic chest conditions)	- Acute disease : (Adults) 200mg, 3 times daily - Chronic disease : (Adults) 200mg, twice daily - Cystic fibrosis : (over 6 years) 200mg, 3 times daily	30C 500C

Respiratory system agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Hederan Syrup		Ivy leaf 70% ethanol fluid ext. 20mg/mL	Catarrhal bronchitis and chronic inflammatory bronchial disease	- Adults and children over 15 years : 10mg, 3 times daily - Children 4~14 years : 100mg, twice daily - Children 2~3 years : 50mg, 3 times daily	500mL
Newkamin Syrup		Pelargonium sidoides 11% ethanol ext.(1~8~10) · glycerin mixed solution(8:2) 0.1716g/mL	Acute bronchitis	- Adults and children Over 12 years : 9mL, 3 times daily - Children 6~12 years : 6mL, 3 times daily - Children 1~6 years : 3mL, 3 times daily	500mL
Levotizine Tab.		Levocetirizine HCl 5mg	Seasonal or perennial allergic rhinitis, chronic idiopathic urticaria, dermatitis and eczema with itching	- Adults and children over 6 years : 1 tablet, once daily regardless of meals - Dose up/down appropriately by age and symptoms	30T 300T
Azelas Tab.		Azelastine HCl 1mg	1. Allergic rhinitis 2. Perennial allergic rhinitis 3. Ocular pruritus associated with allergic conjunctivitis	- 1 tablet, twice daily - Bronchial asthma : 2mg, twice daily (after breakfast and before bedtime)	30T 300T
Montest Tab, 10mg		Montelukast 10mg	1. Asthma (treatment and prevention) 2. Allergic rhinitis	Adults and Adolescents over 15 years : 1 tablet, once daily	28T
Montest Chewable Tab, 4mg		Montelukast 4mg		Children 2~5 years : 1 tablet, once daily	28T
Montest Chewable Tab, 5mg		Montelukast 5mg		Children 6~14 years : 1 tablet, once daily	28T
Momental nasal Spray		Mometasone furoate 50mcg/1dose	1. Seasonal allergic rhinitis (prevention and treatment) 2. Perennial allergic rhinitis 3. Preventive therapy can be 2~4 weeks before planned date of pollen period 4. Nasal polyp in the adults over 18 years	- Children over 12 years and adults : spray twice per each nostril once daily - Children 2~11 years : spray once per nostril once daily	140 dose

Anti-infective agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Amoctam Tab, 500mg		Amoxicillin 250mg Pivoxyl sulbactam 250mg	Otitis media,sinusitis, laryngopharyngitis, bronchitis, pneumonia, upper and lower urethritis, wound infections	1~2 tablets, 3 times daily	30T 300T
Amokle Tab, 375mg		Amoxicillin hydrate 250mg Potassium clavulanate diluted 212,73mg (As Potassium clavulanate 125mg)	Acute/chronic bronchitis, lobar and bronchial pneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, urethritis, pyelonephritis, septic abortion, puerperal period sepsis, pelvic infections, chancroid, gonorrhea, furunculous and abscess, cellulitis, wound infections, intraperitoneal sepsis, peritonitis, osteomyelitis, sepsis, dental infections	Adults and children over 12 years or 40kg of body weight : 250mg as amoxicillin 3 times daily, dose up to 500mg as amoxicillin in case of severe infections	80T
Amokle Tab, 625mg		Amoxicillin hydrate 500mg Potassium clavulanate diluted 212,73mg (As Potassium clavulanate 125mg)			40T
Amokle Syrup		Amoxicillin 40mg/mL Clavulanate potassium 5.7mg/mL		- Mild to moderate infections : 0,625mL/kg - Severe infections : 1,125mL/kg daily in 2 divided doses	50mL
Clamasin Tab.		Clarithromycin 250mg	Lower respiratory tract infection (LTRI), Upper respiratory tract infection (URTI), Mycobacterium infection by mycobacterium avium or mycobacterium intracelluar, eradication of helicobacter pylori in duodenal ulcer patients	- Adults : 250mg, twice daily - Eradication of helicobacter pylori : 500mg, 3 times daily for 14 days	30T 100T
Clamasin Tab, 500mg		Clarithromycin 500mg			30T 200T

Anti-infective agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Roxitirocin Tab.		Roxithromycin 150mg	Laryngopharyngitis, acute bronchitis, tonsillitis, bacterial pneumonia, mycoplasma pneumonia, genital infection, venereal disease excluding gonococcal urethritis, furunculous and abscess, otitis media, sinusitis, folliculitis, boil, carbuncle, lymphangitis (lymphadenitis), whitlow, purulent, perionychia, subcutaneous abscess	- 1 tablet, twice daily - Pneumonia : 2 tablets, once daily before breakfast	30T 500T
Cefler Cap.		Cefaclor 250mg	Otitis media, pneumonia, laryngopharyngitis, tonsillitis, bronchitis, pyelonephritis, cystitis gonococcal urethritis, turuncle, carbuncle, folliculitis, cellulitis, infectious atheroma, subcutaneous abscess, whitlow	- Adults : 250mg, every 8 hours - Children : 20mg/kg, every 8 hours	30C 500C
Saldin Cap, 500mg		Cephradine 500mg	Tonsillitis, pharyngitis, lobar pneumonia, bronchitis, prostatitis, cystitis, pyelonephritis, urethritis, abscess, cellulitis, furunculous, impetigo, otitis media	250~500mg, 4 times daily or 500mg~1g, twice daily	30C 500C
Exen Tab.		Ofloxacin 100mg	1. Susceptible microorganisms : Staphylococcus aureus, Streptococcus pneumoniae, Streptococcus pyogenes, Citrobacter sp., Enterobacter sp., Acinetobacter sp., Serratia sp., E. coli, H. influenzae, Klebsiella pneumoniae, Neisseria gonorrhoeae, Proteus mirabilis, Pseudomonas aeruginosa, Chlamydia trachomatis 2. Indication : Mild to moderate infection caused by susceptible strains of the designated miroorganisms.	3~6 tablets, 2~3 divided doses	30T 500T
Eprosin Tab.		Ciprofloxacin HCl 291mg (As Ciprofloxacin 250mg)	1. Infection disease by E. coli, Shigella, Salmobella, Citrobacter, Klebsiella, Enterobacter, Serratia, Hafnia, Proteus, Pseudomonas, Neisseria, Acinetobacter, Streptococcus, Chlamydia, Staphylococcus, Corynebacterium, Bacteroides, Clostridium 2. Infection : respiratory infection, ear · nose · throat infection, infection of oral · tooth · jaw, infection of kidney or urinary tract, genital infection including gonorrhea, GIT infection, bile secretory duct infection, infection and injury of soft tissue, bone and joint infection, infection of gynecology and obstetrics, septicemia, meningitis, peritonitis, ophthalmology infection	1~2 tablets, twice daily between meals	30T 500T
Levicin Tab.		Levofloxacin 100mg	Infection of the respiratory tract, skin, soft tissue and urinary tract	1 tablet, 2~3 times daily	30T 200T

Antipyretic, Analgesic, Anti-inflammatory agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Loxpain Tab.		Loxoprofen sodium hydrate 68,1mg (As Loxoprofen sodium 60mg)	Rheumatoid arthritis, osteoarthritis (degenerative arthritis), ankylosing apodylitis, tenonitis, arthritis, acute gout and dysmenorrhoea	1 tablet, 3 times daily	30T 300T
Mupain Tab.		Acetaminophen 325mg Tramadol HCl 37,5mg	Moderate to severe pain	2 tablets, every 6 hours	30T 300T
Mupain Semi Tab.		Acetaminophen 162,5mg Tramadol HCl 18,75mg		4 tablets, every 6 hours	30T 300T
Mupain ER Tab.		Acetaminophen 650mg (IR 325mg, ER 325mg) Tramadol HCl 75mg		1 tablet, every 12 hours	30T 100T
Mupain Semi ER Tab.		Acetaminophen 325mg (IR 162,5mg, ER 162,5mg) Tramadol HCl 37,5mg		2 tablets, every 12 hours	30T 100T

Antipyretic, Analgesic, Anti-inflammatory agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Aceline Tab.		Acetoclofenac 100mg	1. Inflammation and pain associated with toothache and trauma, back, sciatica, pain associated with postoperative, Perineal lateralis section and postrartum and non-articular rheumatism 2. Rheumatoid arthritis, ankylosing spondylitis, osteoarthritis, scapulohumeral periarthritis.	1 tablet, twice daily	30T 500T
Moruming Tab.		Morniflumate 350mg	1. Respiratory infection : laryngopharyngitis, trachetis, bronchitis 2. Ear · nose · throat infection : pansinusitid, otitis, tonsillitis, pharyngitis 3. Osteoarthritis	1 tablet, twice daily	30T 300T
Melokcam Cap.		Meloxicam 7.5mg	Rheumatoid arthritis, osteoarthritis, ankylosing spondylitis daily	- Ostarthritis : 1 capsule, once daily - Rheumatoid arthritis, ankylosing spondylitis : 2 capsules, once daily	30C 500C

Antifungal agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Fuzole Cap.		Fluconazole 50mg	1. Vaginal candidiasis 2. Oropharyngeal and esophageal candidiasis 3. Candida urinary tract infection, peritonitis, and systemic, Candida infection including candidemia, disseminated candidiasis, and pneumonia 4. Cryptococcal meningitis, Before prescribing fluconazole for AIDS patients with cryptococcal meningitis	- Vaginal candidiasis : 3 capsules, once daily - Oropharyngeal and esophageal candidiasis : 1 capsule, once daily for 7-14 days - Onychomycosis : 3 capsules, once weekly	30C 100C
Icozole Tab.		Itraconazole 100mg	1. Vaginal candidiasis 2. Pityriasis versicolor 3. Mucocutaneous candidiasis, 4. Oropharyngeal and esophageal candidiasis 5. Onychomycosis due to dermatophytes (tinea unguium) 6.Systemic fungal infections : aspergillosis, candidiasis, cryptococcosis, paracoccidioidomycosis	- Vaginal candidiasis : 2 tablets, twice daily - Pityriasis versicolor : 2 tablets, once daily for 7 days - Oropharyngeal candidiasis : 1 tablet, once daily for 15 days - Tinea corporis, tenia cruris : 1 tablet, once daily for 15 days - Tenia manus, tenia penis : 1 tablet, once daily for 15 days - Onychomycosis : 2 tablet, twice daily for 3 months	28T 100T

Antiviral agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Famdiac Tab.		Famciclovir 250mg	1. Herpes zoster 2. Genital herpes infection and inhibition of recurrent herpes simplex	- Herpes zoster infection : 1 tablet, 3 times daily for 7 days - Recurrent gonococcal infection : 1/2 tablet, twice daily for 5 days	21T

Anticancer

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Iscador M 0.01mg		Viscum Album (Mali) Soluble Extract (1-5) 0.05mg/mL	1. Cancer 2. Prophylaxis of recurrent cancer after surgery 3. Malignant disease of precancerous lesion in hemopoietic organ 4. Stimulation of marrow function	- Initial stage : Increase from series 0 step by step, 2-3 times a week by subcutaneous injection - Maintenance stage : Control individually optimized dose to maintain	7A

Anticancer

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Iscador M 0.1mg		Viscum Album (Mali) Soluble Extract (1-5) 0.5mg/mL	1. Cancer 2. Prophylaxis of recurrent cancer after surgery 3. Malignant disease of precancerous lesion in hemopoietic organ 4. Stimulation of marrow function	- Initial stage : Increase from series 0 step by step, 2-3 times a week by subcutaneous injection - Maintenance stage : Control individually optimized dose to maintain	7A
Iscador M 1 mg		Viscum Album (Mali) Soluble Extract (1-5) 5mg/mL			7A
Iscador M 10mg		Viscum Album (Mali) Soluble Extract (1-5) 50mg/mL			7A
Iscador M 20mg		Viscum Album (Mali) Soluble Extract (1-5) 100mg/mL			7A
Iscador Q 0.01mg		Viscum Album (Quercus) Soluble Extract (1-5) 0.05mg/mL			7A
Iscador Q 0.1mg		Viscum Album (Quercus) Soluble Extract (1-5) 0.5mg/mL			7A
Iscador Q 1mg		Viscum Album (Quercus) Soluble Extract (1-5) 5mg/mL			7A
Iscador Q 10mg		Viscum Album (Quercus) Soluble Extract (1-5) 50mg/mL			7A
Iscador Q 20mg		Viscum Album (Quercus) Soluble Extract (1-5) 100mg/mL			7A

Dermatologic agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Dupina Tab.		Finasteride 1mg	Male alopecia (18-41 years)	1 tablet, once daily	30T
Acnon Soft Cap.		Isotretinoin 10mg	Severe recalcitrant cystic acne vulgaris (nodular acne)	- Initial dose : 0.5mg/kg daily, depending on the symptoms - Maintenance dose : 0.5~1.0mg/kg daily, severe acne patients is 2mg/kg daily - Low dose is once daily and high dose is twice daily with meals	30C 100C

Urogenital system agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Tetrin Tab, 2mg		Terazosin HCl 2,347mg (As Terazosin 2mg)	1. Dysuria due to benign prostate hyperplasia (BPH), neurogenic bladder 2. Hypertension (mild to moderate)	- Dysuria due to BPH ·Initial dose : 1/2 tablet before bedtime ·Maintenance dose : 5~10mg, once daily - Hypertension ·Initial dose : 1/2 tablet before bedtime ·Maintenance dose : 2~10mg, once daily	30T 100T
Tarocin SR Tab, 0.2mg		Tamsulosin HCl 0.2mg	1. Dysuria due to benign prostate hyperplasia (BPH) 2. Neurogenic bladder	1 tablet, once daily	30T 100T
Enpisca Tab,		Finasteride 5mg	Benign prostate hyperplasia (BPH)	1 tablet, once daily	30T
Urofix Tab,		Propiverine HCl 20mg	1. Urinary incontinence (urgency and frequency in unstable bladder conditions) 2. Idiopathic bladder instability 3. Neurogenic bladder (detrusor hyperreflexia)	1 tablet, once daily	30T 100T

Central Nervous System agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Depren Cap,		Fluoxetine HCl 22.4mg (As Fluoxetine 20mg)	Depression, obsessive-compulsive disorder (OCD), agoraphobia, bulimia nervosa, premenstrual dysphoric disorder (PMDD)	20~80mg, a daily	30C 300C
Depren Cap, 10mg		Fluoxetine HCl 11.2mg (As Fluoxetine 10mg)			30C 300C
Bupron Tab,		Bupropion HCl 100mg	Depression	- Initial dose : 1 tablet, twice daily - After 4 days : 1 tablet, 3 times daily	100T

Antiobesity agent

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Fastin Tab,		Phentermine HCl 37.5mg (As Phentermine 30mg)	Short-term treatment of obesity, used as an adjunct to behavior modifications including dietary modification and appropriate exercise	1 tablet, once daily	30T 100T
PTM Tab,		Phentermine HCl 18.75mg (As Phentermine 15mg)		2 tablets, once daily or 1 tablet, 1-2 daily	30T 100T
Phentin Tab,		Phendimetrazine tartrate 35mg		1 tablet, 2-3 times daily	300T
Mermaid Tab,		Alginate acid 200mg Carboxymethylcellulose sodium 100mg	Adjuvant therapy for weight loss by reduction of food intake	2 tablets, 3 times daily	300T
Zerodown Cap, 60mg		Orlistat 60mg	Obesity (including weight loss and weight maintenance) in conjunction with a reduced-calorie diet and appropriate exercise	1 capsule, 3 times per day with each main meal containing fat, taken during the meal or up to 1 hour after the meal	84C
Zerofat Cap,		Orlistat 120mg		1 capsule, 3 times per day with each main meal containing fat, taken during the meal or up to 1 hour after the meal	84C

Others

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Narex Tab,		Naltrexone HCl 25mg	Alcohol dependency combined with appropriated management program and prevention blocking effect of exogenic opioid	2 tablets, once daily	30T 100T
Topin Tab, 25mg		Topiramate 25mg	Prevention of epilepsy, migraine	- Epilepsy : 1 tablet in the evening during the first 1 week, and increase 25 to 50mg per day at the interval of 1~2 weeks, twice daily (recommended dosage as single therapy is 100~200mg per day) - Prevention of migraine : 1 tablet in the evening during the first 1 week, and increase by 25mg at the interval of 1 week	30T 100T
Topit Tab,		Topiramate 100mg			30T 100T
Calurin S Tab,		Ephedrine HCl 25mg Caffeine anhydrous 25mg Acetaminophen 200mg	Mitigation of common cold symptoms (faucal (throat), coughing, phlegm, chills, fever, headache, arthralgia, muscle pain)	1 tablet, 3 times daily	300T
Naomi Tab,		Banggi/Hwanggiang Dried Ext. (7.52→1) 244mg	Obesity with pale skin tone, frequent fatigue and sweating (poor elasticity in the muscle), arthralgia, edema	4 tablets, 3 times daily	360T
Histan F Tab, 16mg		Betahistine HCl 16mg	Dizziness, tinnitus, hearing loss by Meniere's disease	- Initial dose : 16mg, 3 times daily - Maintenance dose : 24~48mg, a daily	30T 300T
Histan F Tab, 24mg		Betahistine HCl 24mg			30T 300T
Chol-Down Soft Cap.,		Pantethine 62.5mg Unsaponifiable Matter of soybean oil 100mg Vitamine E(Tocopherol) 22.67 I.U.	1. Improvement of hypercholesteremia 2. Mitigation of peripheral circulatory disorder (cold hands and feet, tingling) with hypercholesteremia	Adults : 2 capsules, 3 times daily after meals	300C
Cholesten Soft Cap,		Safflower oil 333.33mg (As linol 250mg) Tocopherol acetate 6.67mg Pyridoxine HCl 0.67mg	Improvement of lipid metabolism by obesity, diabetic, primary hypercholesteremia	3 capsules, 3 times daily (30 minutes before meals)	300C
Cambel Tab,		Bangpoongtongsungsan Dried Ext. (4.64→1) 237.5mg	Following symptoms of patients with much abdominal subcutaneous fat and constipation: symptomatology of hypertension (palpitation, shoulder discomfort, flush), obesity, swelling, constipation	2 tablets, 3 times daily before or between meals	500T
D-catin Tab,		L-carnitine 330mg	1. Primary and secondary carnitine deficiency 2. Myocardial metabolism disease by ischemic heart disease : angina, acute myocardial infarction	2~3g daily in 2-3 divided doses	90T
Mano-L Tab,		Aloe extract 100mg Magnesium hydroxide 500mg Ursodesoxycholic acid 10mg	Mitigation of constipation and the following symptoms by constipation: Anorexia (anepithymia), bloating, intestinal abnormal fermentation, hemorrhoids	- Adults and children over 15 years : 1 capsule, 3 times daily - Children 11-14 years : 1 capsule, twice daily before (between or after) meals	300T
Daien Cap,		Green tea powder 250mg Orthosiphon powder 150mg	Complementation of weight loss in obesity or over weight	1~2 capsules, 3 times daily	300C
Vita B Tab,		Folic acid 250µg Cyanocobalamin 500µg Pyridoxine HCl 50mg	1. Supply of vitamin B ₆ for the following - Pregnancy · breastfeeding period - Reduced physical strength during or after sickness 2. Mitigation of the following symptoms : commissural cheilitis (angular stomatitis), cheilitis (cheilitis grandularis), stomatitis (canker sore), glossitis (aphthous ulcer), eczema, dermatitis	Adults and children over 8 years : 1 tablet, twice daily	300T
Tiparon Tab,		Tiropamide HCl 100mg	Acute convulsive pain (hepatobiliary colic, abdominal colic by reasons), abdominal convulsion and pain (renal colic, gastrointestinal dyskinesia syndrome, cholelithiasis, cholecystitis, pain due to postoperative synechia)	1 tablet, 2-3 times daily	30T 500T
Citracal F Tab,		Calcium citrate 750mg (As Calcium 0.158g) Cholecalciferol 4mg (As Vitamin D 400I.U)	Calcium deficiency and supply of calcium, poor growth of bone and teeth, prevention of rickets and supply of vitamin D	2 tablets, once daily	300T

PART2 | Injections

Antipyretic, Analgesic, Anti-inflammatory

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Cnac Inj. 2mL		Diclofenac β--dimethylamino ethanol 90mg/2mL	Rheumatoid arthritis, osteo arthritis, ankylosing spondylitis, inflammation, pain after surgery and injury, acute gout, nephritic and hepatic colic	1 ampule a day, inject deeply on the upper hip	50A

Cardiovascular

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Gitako Inj.		Ginkgo leaf extract 17.5mg/5mL	Brain blood circulatory disorder and brain nutrition disorder (cerebrovascular insufficiency), decline of intellectual and awareness ability, brain function disorder with dizziness, tinnitus	Inject intramuscularly or intravenously 5~10mL a day or inject intravenously 15mL every 2 days	25A
Mesocartin Inj.		L--carnitine 2g/10mL	1. Primary, secondary carnitine deficiency 2. Myocardial metabolism disorder by ischemic heart disease: angina, acute myocardial infarction 3. Carnitine deficiency in End Stage Renal Disease (ESRD) patients receiving hemodialysis	Primary, secondary carnitine deficiency : Inject intramuscularly or intravenously 2~3g a day in adults throughout 2~3 times	10A
Forcecholine Inj.		Choline alfoscerate 1g/4mL	1. Secondary symptom and metamorphosis or degenerative brain-organic psychiatric syndrome by cerebrovascular deficiency: reduced memory, derangement, disorientation by reduced willingness and spontaneity, reduced willingness and spontaneity, reduced concentration 2. Emotional and behavioral change: emotional insecurity, sensitive to stimulation, indifferent to surrounding 3. Senile pseudodepression	Inject intramuscularly or intravenously 1g per injection daily	10A

Anticonvulsive

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Mine-M Inj.		Magnesium sulfate 0.5g/5mL	Convulsion, eclampsia, electrolyte supply (hypomagnesemia), Spasticity of the uterus (parturifacient)	Slowly inject intramuscularly or intravenously 1~5g, very slowly inject intramuscularly or intravenously 2~4g throughout several times a day for electrolyte supply	25A

Vitamins

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Multivita Inj.		Cyanocobalamin 5μg Biotin 60μg Folic acid 400μg Tocopherol acetate 10mg Dexpanthenol 15mg Nicotinic acid amide 40mg Ascorbic acid 100mg Riboflavin sodium phosphate 3.6mg Thiamine HCl 3.81mg Pyridoxine HCl 4.86mg Ergocalciferol 5μg(200IU) Retinol palmitate 1.815mg(3,300IU) /1mL	1. Maintenance therapy of the vitamins for patient who nutrients are supplied with fluid 2. Prevention of vitamins deficiency for the following diseases (excluding vitamin K) – Surgical surgery, severe burn, fracture and other injuries – Severe infection, coma	Adults and children over 11 years : after melting this medicine by putting it into 5mL distilled water for injection by gently shaking, mix it in 500~1,000mL of glucose injection solution, saline or equivalent injection solution for intravenous injection within 5 minutes, and inject by intravenous infusion once daily	10V

Vitamins

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Pentavita Inj.		Dexpanthenol 500mg/2mL	Pantothenic acid deficiency (treatment and prevention), supply for increased demand of pantothenic acid and insufficient intake of pantothenic acid from meals	Inject intramuscularly or intravenously 20~100mg per injection 1~2 times daily, for postoperative paresis of intestine, inject 50~500mg 1~3 times daily, if necessary increase up to 6 times	50A
Furmine Inj. 5mg		Fursultamine HCl 54.6mg/10mL	Vitamin B1 deficiency (treatment and prevention), supply for increased demand of vitamin B1 and insufficient intake of vitamin B1 from meals	Adults : Inject on the subcutaneous muscle or intravenously 5~100mg daily	10A
Hexavita Inj.		Pyridoxine HCl 100mg/2mL	Vitamin B6 deficiency (treatment and prevention), supply for increased demand of vitamin B6 and insufficient intake of vitamin B6 from meals	Inject on the subcutaneous muscle or intravenously 10~100mg throughout 1~2 times daily	50A
Vitamin C Injection 10g		DaehanNewpharm Ascorbic acid 10g/20mL	Vitamin C deficiency (treatment and prevention), in case of increased demand of vitamin C	Inject on the subcutaneous muscle or intravenously 50~1,000mg throughout once daily or several times a day	10V
Vita-D Inj.		Cholecalciferol 5mg (200,000IU)/1mL	vitamin D deficiency (treatment and prevention)	- Adults : Inject intramuscularly 200,000IU - Adolescents : Inject intramuscularly 200,000IU - Elderly people : Inject intramuscularly 100,000IU	5A

Inorganic

Product name	Icon	Ingredient Content	Indication	Dosage	Package
MultiTrace Inj.		Selenious acid 327μg Chromic chloride hydrate 205μg Cupric sulfate pentahydrate 15.7mg Manganese sulfate monohydrate 3.08mg Zinc sulfate heptahydrate 43.9mg /10mL	Supply of trace element (zinc, copper, mangan, chrome) in the form of intrajugular nutrition supply	Use by doctor's prescription : however, confirm the microelement amount in plasma to control dose	10V
Selen Inj.		Selenious acid 654μg/10mL	Selenium supplementation in total parenteral nutrition (TPN)	Metabolically stable adults: inject 20~40 μg of selenium a day during TPN this drug includes 40 μg/mL of selenium)	10V
DHNP Magnesium Chloride Inj. 10%		Magnesium chloride 1g/10mL	1. Torsade de pointe 2. Acute hypokalemia accompanying hypomagnesemia 3. Magnesium supplementation for unbalanced electrolyte 4. Magnesium supplementation in TPN 5.Preeclampsia (gestosis) (treatment and prevention)	Torsade de pointe : slowly inject intravenously 8mmol of magnesium (1.6g of magnesium chloride)	10A
Zinc Inj. 5mL		Zinc sulfate hydrate 22mg (As Zinc 5mg)	Zinc supplementation in TPN	Metabolically stable TPN patients: 2.5~4mg daily	10V
Zinc Inj. 10mL		Zinc sulfate hydrate 44mg (As Zinc 10mg)			10V

Amino acid

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Placein M Inj.		Placenta extract 2mL/2mL	Improvement of menopausal disorder symptoms	Inject hypodermically 2mL per injection, 3 times a week for 2 weeks (total 6 times)	10A

Non Steroidal Anti-inflammatory

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Bicolamin Inj.		Hydroxocobalamin 5mg/2mL	Acute neuritis, multiple neuritis, sciatic neuralgia, rheumatoid cervical syndrome, radiculalgia, adhesive capsulitis, gastritis, neuralgia by cancer or injury, neuralgia by shingles and migraine	Inject intramuscularly or intravenously once daily	50A

Others

Product name	Icon	Ingredient Content	Indication	Dosage	Package
ENG Inj.		Adenosine triphosphate disodium trihydrate 21.96mg/2mL (As Adenosine triphosphate disodium 20mg)	Improvement of hepatic function in myasthenia, heart failure, chronic infection after by head injury, accommodative asthenopia	Inject hypodermically, intramuscularly or intravenously 5~20mg per injection in normal adults 1~2 times daily	10A
Lycoris Inj.		Ammonium Glycyrrhizinate 26.5mg Glycine 200mg Cysteine Hydrochloride Monohydrate 10mg/10mL	Complementation of rashes, eczema, allergic skin trouble, addiction, improvement of hepatic functions in chronic liver disease	Adults : Inject intravenously or through intravenous drip 2~20mL per injection, once daily, for chronic liver disease, inject 40~60mL per injection once daily. In case of increasing dose, do not exceed maximum 100mL a day	50A
Cindella Inj.		Thioctic acid 25mg/5mL	Supply for increased demand of thioctic acid (intensive physical labor), Leigh syndrome (subacute necrotic encephalomyelitis), addictive (by streptomycin and kanamycin) and noise induced (occupational) internal ear deafness	Inject 10~25mg intravenously in adults once daily	10A
Lipoic Inj.		Thioctic acid tromethamine 476.4mg/12mL (As Thioctic acid 300mg)	Mitigation of diabetic polyneuropathy	For severe case, inject intravenously 600mg of thioctic acid daily for 2~4 weeks, and afterwards 600mg daily	10A
Bione Inj.		Glutathione(reduced) 600mg	Prevention of neurotic disease by cisplatin or similar chemical category therapy	For severe case, inject intramuscularly or through intravenous drip 600~1,200mg daily	10V
Luthione Inj.		Glutathione(reduced) 600mg			10V
Luthione Inj. 1200mg		Glutathione(reduced) 1200mg			10V
New-DN Inj.		Polydeoxyribonucleotide sodium 5.625mg/3mL	Wound by skin graft and tissue restoration	Inject intramuscularly or hypodermically 1 vial daily	10V
Liporase Inj.		Hyaluronidase 1,500 I.U.	Increase of permeability in hypodermic injection or intramuscular injection, local anesthetic and hypodermic infusion, resorption of excessive body fluid and blood existing excessively in a tissue	During hypodermic infusion (hypodermoclysis): Melt 1 vial into 1mL of injection solution or saline, inject to a relevant area before hypodermic infusion	10V

Health supplement

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Ascorbic acid 3g		Ascorbic acid 3g	Supply of ascorbic acid	Take with water or beverage	100 Pouches
Ultra fine ascorbic acid		Ascorbic acid 3g	Supply of ascorbic acid	Take with water or beverage	90 Pouches
MSM		MSM(Methyl sulfonylmethane) 2g	Assistance for joint and cartilage health	Take 1 pack (2g) with water, once daily	120 Pouches
Super MSM Protocol		MSM(Methyl sulfonylmethane) 750mg Vitamin C 300mg	Assistance for joint and cartilage health, effective relief of neuralgia, removal of extensive free oxygen radicals, supply of ascorbic acid	2 tablets, once daily	180T
Aniplus Cap.		Vitamin C, Magnesium oxide, Crystalline cellulose, L-Tyrosine	[Vitamin C] Necessary for the formation of connective tissue and function maintenance, necessary for iron absorption, necessary to protect cells from harmful oxygen [Magnesium] Necessary for energy use, necessary for nerve and muscle functions	1 capsule, BID	120C
Carboglu-X Cap.		Vitamin B ₁ Nitrate, Vitamin B ₂ , Vitamin B ₆ HCL, Vitamin C, Crystalline cellulose, White Kidney Bean extract, Citrus Unshiu Peel extract, Milk calcium	[Vitamin B ₁] Necessary for carbohydrate and energy metabolism [Vitamin B ₂] Necessary for creating energy in the body [Vitamin B ₆] Necessary for using protein and amino acid, necessary to maintain the normal homocysteine level in blood [Vitamin C] Necessary for the formation of connective tissue and function maintenance, necessary for iron absorption, necessary to protect cells from harmful oxygen	2 capsules, twice daily	300C

Medical device

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Lafillere MAX		Hyaluronic acid 20mg	Temporary improvement of facial wrinkles	Inject into the dermal layer of facial wrinkles	1,1 mL/s
SSONDAR M9 Injector		- 9-Pin Needle - Needle Size: 32G	Injector exclusively for Hyaluronic acid injection		1 Set

Other drug

Product name	Icon	Ingredient Content	Indication	Dosage	Package
Lidocan Cream		Lidocaine 96mg/g	Reduction of tactile sensitivity of the penis	As ointment, apply 0.25~0.66g (as lidocaine, apply 23~62mg)	500g

New Platform for Life Care



Headoffice (R & D) : 66 Pharmaceutical Industries Complex 1-gil Hyangnam-eup, Hwasung-si, Gyeonggi-do, Korea
Seouloffice : 61 Namhyeon 3-gil, Gwanak-gu, Seoul, Korea, 08806
TEL : +82 2 581 2333