

# PRODUCT LIST | HISTORY

- 1984 | Establishment of Daehan Sinyak
  - | Received approval for animal drug manufacturing
- 1994 | Acquired a pharmaceutical company
- 2001 | Changed company name to Dae Han New Pharm
  - | Received KVGMP certification for animal drugs
- 2002 | Initial Public Offering (IPO)
- 2005 | Awarded for outstanding export performance (USD 3M)
- 2007 | Awarded for outstanding export performance (USD 5M)
- 2015 | Awarded for outstanding export performance (USD 10M)
- 2021 | Completion of R&D center in Pangyo
- 2022 | Changed company name to Daehan Nupharm
- 2023 | Expansion of Hyangnam factory

# PRODUCT LIST | C O L O R S

## ATC code | 1st level

The 1st level is the main classification based on the anatomical site where the medicine acts.

There are 14 levels from A to V.

<b>A</b>	Alimentary tract and metabolism	<b>L</b>	Antineoplastic and immunomodulating agents
<b>B</b>	Blood and blood forming organs	<b>M</b>	Musculo-skeletal system
<b>C</b>	Cardiovascular system	<b>N</b>	Nervous system
<b>D</b>	Dermatologicals	<b>P</b>	Antiparasitic products, insecticides and repellents
<b>G</b>	Genito-urinary system and sex hormones	<b>R</b>	Respiratory system
<b>H</b>	Systemic hormonal preparations, excluding sex hormones and insulins	<b>S</b>	Sensory organs
<b>J</b>	Antiinfectives for systemic use	<b>V</b>	Various

## ATC code | 2nd level

The 2nd level is a detailed classification based on therapeutic/pharmacological subgroups, and is denoted by numbers from 1 to 16.

<b>01</b>	<b>02</b>	<b>03</b>	<b>04</b>	<b>05</b>	<b>06</b>	<b>07</b>	<b>08</b>
<b>09</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>

# Circulatory System











C

08

09

10







Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Avatri Tab. 5/80/10mg</b> Amlodipine besylate 6.94mg (as amlodipine 5mg) / Valsartan 80mg / Atorvastatin calcium hydrate 10.85mg (as atorvastatin 10mg)	<ul style="list-style-type: none"> <li>* Intended for patients requiring simultaneous administration of the amlodipine/valsartan combination and atorvastatin.</li> <li>* For essential hypertension where blood pressure is inadequately controlled with either amlodipine or valsartan monotherapy.</li> <li>* Atorvastatin               <ol style="list-style-type: none"> <li>1. To reduce the risk of cardiovascular diseases.</li> <li>2. For the management of dyslipidemia.</li> </ol> </li> </ul>	Take 1 tablet once daily, preferably at the same time each day.	30T/Bottle	C10BX
	<b>Avatri Tab. 5/80/20mg</b> Amlodipine besylate 6.94mg (as amlodipine 5mg) / Valsartan 80mg / Atorvastatin calcium hydrate 21.7mg (as atorvastatin 20mg)				
	<b>Avatri Tab. 5/160/10mg</b> Amlodipine besylate 6.94mg (as amlodipine 5mg) / Valsartan 160mg / Atorvastatin calcium hydrate 10.85mg (as atorvastatin 10mg)				
	<b>Avatri Tab. 5/160/20mg</b> Amlodipine besylate 6.94mg (as amlodipine 5mg) / Valsartan 160mg / Atorvastatin calcium hydrate 21.7mg (as atorvastatin 20mg)				
	<b>Amloke Tab.</b> Amlodipine besylate 6.944mg (as amlodipine 5mg)	<ul style="list-style-type: none"> <li>* Myocardial ischemia caused by hypertension, stable angina, or coronary vasospasm and vasoconstriction (variant angina)</li> <li>* Patients recently diagnosed with coronary heart disease through angiography, without heart failure, or with a cardiac output above 40%:               <ol style="list-style-type: none"> <li>1. To reduce the risk of hospitalization due to angina.</li> <li>2. To lower the risk of coronary revascularization.</li> </ol> </li> </ul>	<ul style="list-style-type: none"> <li>* Take 5mg orally once daily.</li> <li>* The dose may be increased to a maximum of 10mg per day, adjusted based on age and symptoms.</li> </ul>	30T/Bottle 300T/Bottle	C08CA01
	<b>Lozantan Plus Tab.</b> Losartan potassium 50mg / Hydrochlorothiazide 12.5mg	Hypertension inadequately controlled by monotherapy	<ul style="list-style-type: none"> <li>* Take 1 tablet once daily.</li> <li>* Can be administered with or without food.</li> </ul>	30T/Bottle 300T/Bottle	C09DA01

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>NP Forge Tab.</b> 5/80mg Amlodipine 5mg / Valsartan 80mg				
	<b>NP Forge Tab.</b> 5/160mg Amlodipine 5mg / Valsartan 160mg	Essential hypertension inadequately controlled by either amlodipine or valsartan monotherapy	* Take 1 tablet once daily.  * Can be administered with or without food.	30T/Bottle	C09DB01
	<b>NP Forge Tab.</b> 10/160mg Amlodipine 10mg / Valsartan 160mg				
	<b>Tertan Tab.</b> 40mg Telmisartan 40mg	* Essential hypertension	* Essential hypertension: Take 40mg orally once daily. If needed, the dose may be increased to 80mg once daily.	30T (10T/PTP*3) /Pack	C09CA07
	<b>Tertan Tab.</b> 80mg Telmisartan 80mg	* Cardiovascular disease risk reduction	* Cardiovascular disease risk reduction: Take 80mg orally once daily.		
	<b>Enpista Tab.</b> 40/5mg Telmisartan 40mg / Amlodipine 5mg				
	<b>Enpista Tab.</b> 40/10mg Telmisartan 40mg / Amlodipine 10mg	For the treatment of essential hypertension when blood pressure is not adequately controlled with telmisartan or amlodipine monotherapy.	* Take 1 tablet once daily.  * Can be administered with or without food.	30T (10T/PTP*3) /Pack	C09DB04
	<b>Enpista Tab.</b> 80/5mg Telmisartan 80mg / Amlodipine 5mg				
	<b>Newmesartan Tab.</b> 10mg Olmesartan medoxomil 10mg		* The recommended starting dose is 10mg once daily.  * This medication can be administered with or without food.	30T/Bottle	
		Treatment of essential hypertension	* If blood pressure is not adequately controlled, the dose may be increased to 20mg once daily.		C09CA08
	<b>Newmesartan Tab.</b> 20mg Olmesartan medoxomil 20mg		* If additional blood pressure reduction is required, the dose can be further increased up to a maximum of 40mg per day	30T/Bottle 100T/Bottle	

# Circulatory System

C

09

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Newvikar Tab. 5/20mg</b> Amlodipine besylate 6.944mg (as amlodipine 5mg) / Olmesartan 20mg				
	<b>Newvikar Tab. 5/40mg</b> Amlodipine besylate 6.944mg (as amlodipine 5mg) / Olmesartan 40mg	Treatment of essential hyper- tension when blood pressure is not adequately controlled with amlodipine or olmesartan medoxomil monotherapy.	Take 1 tablet once daily.	30T (10T/PTP*3) /Pack	C09DB02
	<b>Newvikar Tab. 10/40mg</b> Amlodipine besylate 13.888mg (as amlodipine 10mg) / Olmesartan 40mg				
	<b>Newvikar Plus Tab. 5/20/12.5mg</b> Amlodipine besylate 6.944mg (as amlodipine 5mg) / Olmesartan medoxomil 20mg / Hydrochlorothiazide 12.5mg				
	<b>Newvikar Plus Tab. 5/40/12.5mg</b> Amlodipine besylate 6.944mg (as amlodipine 5mg) / Olmesartan medoxomil 40mg / Hydrochlorothiazide 12.5mg	Essential hypertension when blood pressure is not adequately controlled with combination therapy of amlodipine and olmesartan medoxomil.	Take 1 tablet once daily.	28T (14T/PTP*2) /Pack	C09DX03
	<b>Newvikar Plus Tab. 10/40/12.5mg</b> Amlodipine besylate 13.888mg (as amlodipine 10mg) / Olmesartan medoxomil 40mg / Hydrochlorothiazide 12.5mg				


Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Newtor Tab. 10mg</b> Atorvastatin calcium 10.85mg (as atorvastatin 10mg)	* Risk reduction for the following cardiovascular diseases For adult patients with multiple risk factors for coronary heart disease but no clinical evidence of coronary heart disease: 1. Reduction in the risk of myocardial infarction 2. Reduction in the risk of stroke 3. Reduction in the risk of revascularization procedures and chronic stable angina  * Hyperlipidemia and other related conditions	* Take in a dose range of 10mg to 80mg once daily.  * Can be administered at any time of day, with or without food.	30T/Bottle 100T/Bottle	C10AA05
	<b>Newtor Tab. 20mg</b> Atorvastatin calcium 21.69mg (as atorvastatin 20mg)				
	<b>Newtozet Tab. 10/10mg</b> Ezetimibe 10mg / Atorvastatin Calcium Trihydrate 10.85mg (as atorvastatin 10mg)	* Primary hypercholesterolemia  * Homozygous familial hypercholesterolemia (HoFH)	* Take 1 tablet once daily, with or without food.  * Primary hypercholesterolemia: The recommended initial dose is 10/10mg or 10/20mg once daily, with a dosage range of 10/10mg to 10/80mg per day.  * Homozygous familial hypercholesterolemia (HoFH): The recommended dose is 10/40mg or 10/80mg once daily.	30T (10T/PTP*3) /Pack	C10BA05
	<b>Newtozet Tab. 10/20mg</b> Ezetimibe 10mg / Atorvastatin Calcium Trihydrate 21.7mg (as atorvastatin 20mg)				
	<b>Newtomega Soft Cap. 5/1,000mg</b> Atorvastatin Calcium Trihydrate 5.425mg (as Atorvastatin 5mg) / Omega-3-acid Ethyl Esters 90 (1,000mg)	* Treatment of complex (type IIb) dyslipidemia in adult patients at high risk of coronary heart disease (CHD) where: 1. LDL-cholesterol levels are adequately controlled 2. Triglyceride levels remain inadequately controlled despite atorvastatin 20mg monotherapy.	Adults: Take 4 capsules once daily with meals to avoid gastrointestinal discomfort.	80C (8C/Alu-Alu*8) /Pack	C10BA08









# Circulatory System

C

10

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Newsto Tab. 10mg</b> Rosuvastatin calcium 10.4mg (as rosuvastatin 10mg)	* Primary hypercholesterolemia (type IIa, including heterozygous familial hypercholesterolemia) and combined hyperlipidemia (type IIb)	* For primary hypercholesterolemia (type IIa, including heterozygous familial hypercholesterolemia), complex hyperlipidemia (type IIb), primary dysbetalipoproteinemia (type III), and homozygous familial hypercholesterolemia (HoFH): The recommended dose is 10–20mg once daily.	30T/Bottle	C10AA07
	<b>Newsto Tab. 20mg</b> Rosuvastatin calcium 20.8mg (as rosuvastatin 20mg)	* Homozygous familial hypercholesterolemia (HoFH)	* For pediatric patients with heterozygous familial hypercholesterolemia (ages 10 to 17): The recommended dose is 5–20mg once daily.		
	<b>Newbaro Tab. 2mg</b> Pitavastatin calcium 2mg	* Reduces elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglyceride levels in patients with: 1. Primary hypercholesterolemia (heterozygous familial and nonfamilial, Fredrickson type IIa) 2. Mixed dyslipidemia (Fredrickson type IIb) 3. Dietary supplement to increase HDL-cholesterol levels	* Take 1 tablet once daily.  * If LDL-cholesterol lowering effect is insufficient, the dose may be increased up to 4mg per day.	30T/Bottle 100T/Bottle	C10AA08
	<b>Enpitorin Tab. 10/10mg</b> Ezetimibe 10mg / Simvastatin 10mg	* Primary hypercholesterolemia			
	<b>Enpitorin Tab. 10/20mg</b> Ezetimibe 10mg / Simvastatin 20mg	* Homozygous familial hypercholesterolemia (HoFH)	Take 1 tablet once daily in the evening, with or without food.	30T (10T/PTP*3) /Pack	C10BA02



















Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Newpido Tab.</b> Clopidogrel 75mg	<ul style="list-style-type: none"> <li>* Improvement of atherosclerotic symptoms in adults with ischemic stroke, myocardial infarction, or peripheral arterial disease</li> <li>* Improvement of atherosclerotic symptoms in adults with acute coronary syndrome</li> <li>* Reduction of the risk of atherothrombosis and thromboembolism, including stroke, in adults with atrial fibrillation who have one or more vascular risk factors, are not suitable for treatment with vitamin K antagonists, and have a low bleeding risk</li> </ul>	<ul style="list-style-type: none"> <li>* Ischemic stroke, myocardial infarction, peripheral arterial disease: 75mg once daily</li> <li>* Acute coronary syndrome: Loading dose: 300mg once daily / Maintenance dose: 75mg once daily / Take with aspirin 75-325mg once daily.</li> <li>* Atrial fibrillation: 75mg once daily, take with aspirin 75-100mg once daily.</li> </ul>	30T/Bottle 100T/Bottle	<b>B01AC04</b>
	<b>Newpogrel SR Tab.</b> 300mg Sarpogrelate HCl 300mg (IR 100mg, SR 200mg)	Improvement of ischemic symptoms such as ulcers, pain, and a sensation of cold caused by chronic arterial occlusion (e.g., Buerger's disease, arteriosclerosis obliterans, diabetic polyneuropathy, etc.).	<ul style="list-style-type: none"> <li>* Take orally once daily after meals.</li> <li>* Do not chew or crush the tablet; swallow it whole.</li> </ul>	30T/Bottle 100T/Bottle	<b>B01AC</b>
	<b>Cilotal SR Cap.</b> 100mg Cilostazol 100mg	* Improvement of ischemic symptoms such as ulcers, pain, and cold sensation caused by chronic arterial occlusion (e.g., Buerger's disease, arteriosclerosis obliterans, diabetic polyneuropathy)	Take 200mg orally once daily on an empty stomach.	30C/Bottle	<b>B01AC23</b>
	<b>Cilotal SR Cap.</b> 200mg Cilostazol 200mg	* Prevention of recurrence after cerebral infarction (excluding cardiogenic cerebral embolism)			
	<b>Limaglan Tab.</b> Limaprost 5mcg	<ul style="list-style-type: none"> <li>* Improvement of ischemic symptoms such as ulcers, pain, and cold sensation caused by obstructive thromboangiitis (Buerger's disease).</li> <li>* Improvement of subjective symptoms (pain and numbness in the lower extremities) and walking ability in patients with acquired lumbar spinal stenosis (normal results in the SLR test and bilateral intermittent claudication).</li> </ul>	<ul style="list-style-type: none"> <li>* For the improvement of ischemic symptoms such as ulcers, pain, and cold sensation caused by obstructive thromboangiitis (Buerger's disease): Adults are typically administered 2 tablets orally, 3 times a day.</li> <li>* For the improvement of subjective symptoms (pain and numbness in the lower extremities) and walking ability due to acquired lumbar spinal stenosis: Adults are typically administered 1 tablet orally, 3 times a day.</li> </ul>	84T (21T/Alu-Alu*4) /Pack	<b>B01AC28</b>
	<b>Metazin Tab.</b> Trimetazidine HCl 20mg	Combination therapy for symptomatic treatment of stable angina in patients inadequately controlled or intolerant to first-line antianginal therapy	<ul style="list-style-type: none"> <li>* Standard dosage: Take 1 tablet three times daily with meals.</li> <li>* For patients with moderate renal impairment (creatinine clearance 30-60mL/min): Take 1 tablet twice daily, with breakfast and dinner.</li> </ul>	30T/Bottle 300T/Bottle	<b>C01EB15</b>

# Endocrine System

A

10

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>NP Dapa Tab. 10mg</b> Dapagliflozin propanediol hydrate 12.3mg (as dapagliflozin 10mg)	* This medication is used as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.  * Can take as monotherapy or combination therapy.	Take 1 tablet once daily.	28T (14T/PTP*2) /Pack	A10BK01
	<b>Dapaformin SR Tab. 10/500mg</b> Dapagliflozin propane- diol hydrate 12.3mg (as dapagliflozin 10mg) / Metformin HCl 500mg	Take as an adjunct to diet and exercise to improve glycemic control in adult patients with type 2 diabetes mellitus for whom the combined use of dapagliflozin and metformin is appropriate.	Take 1 tablet once daily, preferably with the evening meal.	30T (6T/Alu-Alu*5) /Pack	A10BD15
	<b>Dapaformin SR Tab. 10/1,000mg</b> Dapagliflozin propanediol hydrate 12.3mg (as dapagliflozin 10mg) / Metformin HCl 1,000mg				
	<b>Sitatin Alpha Tab. 50mg</b> Sitagliptin hydrochloride hydrate 59.69mg (as sitagliptin 50mg)	This medication is used as an adjunct to diet and exercise to improve glycemic control in adult patients with type 2 diabetes mellitus when the combined use of sitagliptin and metformin is appropriate.	* Take 1 tablet once daily.	30T (10T/PTP*3) /Pack	A10BH01
	<b>Sitatin Alpha Tab. 100mg</b> Sitagliptin hydrochloride hydrate 113.37mg (as sitagliptin 100mg)		* Maximum daily dose: 100mg		
	<b>Sitafomin Aplha Tab. 50/500mg</b> Sitagliptin hydrochloride hydrate 56.69mg (as sitagliptin 50mg) / Metformin HCl 500mg				
	<b>Sitafomin Aplha Tab. 50/850mg</b> Sitagliptin hydrochloride hydrate 56.69mg (as sitagliptin 50mg) / Metformin HCl 850mg	This medication is used as an adjunct to diet and exercise to improve glycemic control in adult patients with type 2 diabetes mellitus when the combined use of sitagliptin and metformin is appropriate.	* Take twice daily with meals.  * Maximum daily dose: sitagliptin 100mg, metformin 2,000mg	30T (10T/PTP*3) /Pack	A10BD07
	<b>Sitafomin Aplha Tab. 50/1,000mg</b> Sitagliptin hydrochloride hydrate 56.69mg (as sitagliptin 50mg) / Metformin HCl 1,000mg				




Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Sitaformin XR Tab. 50/500mg</b> Sitagliptin hydrochloride hydrate 56.69mg (as sitagliptin 50mg) / Metformin HCl 500mg				
	<b>Sitaformin XR Tab. 50/1,000mg</b> Sitagliptin hydrochloride hydrate 56.69mg (as sitagliptin 50mg) / Metformin HCl 1,000mg	The combination therapy of sitagliptin and metformin is administered as an adjunct to diet and exercise to improve glycemic control in appropriate patients with type 2 diabetes.	* Take with meals, once daily.  * Maximum daily dose: sitagliptin 100mg, metformin 2,000mg	28T/Bottle	A10BD07
	<b>Sitaformin XR Tab. 100/1,000mg</b> Sitagliptin hydrochloride hydrate 113.37mg (as sitagliptin 100mg) / Metformin HCl 1,000mg				
	<b>Duogli Tab.</b> Dapagliflozin citric acid (Micronized) 14.7mg (10mg as dapagliflozin) / Sitagliptin hydrochloride hydrate 113.4mg (100mg as sitagliptin)	The combination therapy of sitagliptin and dapagliflozin is administered as an adjunct to diet and exercise to improve glycemic control in appropriate patients with type 2 diabetes mellitus.	Take 1 tablet once daily, with or without food.	30T (10T/PTP*3) /Pack	A10BD29
	<b>Teneltin Tab. 20mg</b> Teneligliptin hydrochloride hydrate 26mg (as teneligliptin 20mg)	Take as an adjunct to diet and exercise therapy to improve blood sugar control in adult patients with type 2 diabetes.	* Take 1 tablet once daily, with or without food.  * No dosage adjustment is required for patients with renal impairment.  * No dosage adjustment is needed for patients with mild to moderate hepatic impairment.	30T/Bottle 60T/Bottle	A10BH08
	<b>Tenelformin SR Tab. 10/500mg</b> Teneligliptin hydrochloride hydrate 13mg (as teneligliptin 10mg) / Metformin HCl 500mg		* Take 2 tablets once daily with a meal.  * Maximum daily dose: Teneligliptin 20mg, Extended-release Metformin 2,000mg		
	<b>Tenelformin SR Tab. 10/750mg</b> Teneligliptin hydrochloride hydrate 13mg (as teneligliptin 10mg) / Metformin HCl 750mg	Take as an adjunct to diet and exercise therapy to improve blood sugar control in patients with type 2 diabetes for whom the combination of teneligliptin and metformin is appropriate.		30T/Bottle	A10BD28
	<b>Tenelformin SR Tab. 20/1,000mg</b> Teneligliptin hydrochloride hydrate 26mg (as teneligliptin 20mg) / Metformin HCl 1,000mg		Take 1 tablet once daily, with food.		

# Endocrine System


A

10

16

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Glizide Tab.</b> Gliclazide 80mg	Type 2 diabetes (limited to cases where sufficient effect cannot be achieved with diet and exercise alone.)	Adults: Start with 40–80mg per day, take orally once or twice daily.	30T/Bottle 500T/Bottle	<b>A10BB09</b>
	<b>Gerimet Tab.</b> 2/500mg Glimepiride 2mg / Metformin HCl 500mg	* In combination with diet and exercise therapy for patients with non-insulin-dependent diabetes mellitus (type 2): 1. When glycemic control is insufficient with glimepiride or metformin monotherapy. 2. As a replacement for combination therapy with glimepiride and metformin.	Take once or twice daily, just before or with meals.	30T/Bottle 100T/Bottle	<b>A10BD02</b>
	<b>R-foic Tab.</b> R-Thioctic acid tromethamine 480mg	Relief of diabetic polyneuritis	* Take 1 tablet once daily. * Take approximately 30 minutes before breakfast with plenty of water.	30T/Bottle 100T/Bottle	<b>A16AX01</b>


## NEW






	<b>Newlagenta Tab.</b> Linagliptin 5mg	Take as an adjunct to diet and exercise therapy to improve glycemic control in patients with type 2 diabetes.	Take 5mg once daily, regardless of meals.	30T (10T/Alu-Alu*3) /Pack	<b>A10BH05</b>
---	---	---	---	---------------------------------	----------------

# Digestive System

A

02

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
<b>South Korea's First &amp; Only!</b>					
	<b>Rapiduo Tab.</b> 10/350mg Rabeprazole sodium 10mg / Magnesium oxide 350mg	* Gastric ulcer and duodenal ulcer * Erosive or ulcerative gastroesophageal reflux disease (GERD) * Symptomatic relief of gastroesophageal reflux disease (GERD) * Long-term maintenance therapy for gastroesophageal reflux disease (GERD)	Take once daily, 10/350mg	28T (14T/Alu-Alu*2) /Pack	<b>A02BC54</b>

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Rapizole Tab. 10mg</b> Rabeprazole 10mg	<ul style="list-style-type: none"> <li>* Gastric ulcer and duodenal ulcer</li> <li>* Erosive or ulcerative gastroesophageal reflux disease (GERD)</li> <li>* Relief of symptoms of gastroesophageal reflux disease (GERD)</li> <li>* Long-term maintenance therapy for gastroesophageal reflux disease (GERD)</li> </ul>	<ul style="list-style-type: none"> <li>* Gastric and duodenal ulcers: 10–20mg once daily</li> <li>* Erosive or ulcerative GERD: 10–20mg once daily</li> <li>* Symptomatic relief of GERD: 10mg once daily</li> <li>* Long-term maintenance therapy for GERD: 10–20mg per day</li> <li>* Eradication of Helicobacter pylori: 20mg of this agent + 500mg clarithromycin + 1g amoxicillin, twice daily for 7 days</li> <li>* Zollinger-Ellison syndrome: 60–120mg per day; up to 100mg can be administered as a single dose, or 120mg can be divided into two doses of 60mg each.</li> </ul>	28T (28T/PTP*1) /Pack	A02BC04
	<b>Rapizole Tab. 20mg</b> Rabeprazole 20mg	<ul style="list-style-type: none"> <li>* Antibiotic combination therapy for peptic ulcer patients infected with Helicobacter pylori</li> <li>* Zollinger-Ellison syndrome</li> </ul>			
	<b>Omprazole Cap. 20mg</b> Omeprazole 20mg	<ul style="list-style-type: none"> <li>* Short-term treatment of gastric and duodenal ulcers</li> <li>* Treatment of symptoms (e.g., heartburn, vomiting) associated with reflux esophagitis and gastroesophageal reflux disease (GERD)</li> <li>* Maintenance therapy for severe reflux esophagitis or poorly managed peptic ulcers</li> <li>* Zollinger-Ellison syndrome</li> <li>* Antibiotic combination therapy to prevent recurrence of duodenal ulcers caused by Helicobacter pylori infection</li> <li>* Treatment of gastric or duodenal ulcers and erosions caused by nonsteroidal anti-inflammatory drug (NSAID) use</li> <li>* Prevention of gastric and duodenal ulcers, erosions, or dyspepsia in patients on long-term NSAID therapy with a history of gastric or duodenal lesions</li> <li>* Prevention of recurrence of reflux esophagitis</li> </ul>	<ul style="list-style-type: none"> <li>* Take 1 capsule once daily.</li> <li>* Treatment of Zollinger-Ellison syndrome: Take 60–120mg once daily. For doses of 80mg or more per day, divide the total dose into two doses administered twice daily.</li> <li>* Prevention of relapse of gastroesophageal reflux disease: Take 10mg once daily.</li> </ul>	28C/Bottle 300C/Bottle	A02BC01
	<b>SOM Tab. 20mg</b> Esomeprazole 20mg	<ul style="list-style-type: none"> <li>* Gastroesophageal reflux disease (GERD)</li> <li>* Antibiotic combination therapy for eradication of Helicobacter pylori</li> <li>* Short-term therapy for upper gastrointestinal symptoms (pain, discomfort, and burning sensation) associated with nonsteroidal anti-inflammatory drug (NSAID) use (COX-2 non-selective and selective)</li> </ul>	<ul style="list-style-type: none"> <li>* Take 20–40mg once daily</li> <li>* For treatment of Zollinger-Ellison syndrome: 40–80mg twice daily</li> </ul>	28T (14T/Alu-Alu*2) /Pack	A02BC05
	<b>SOM Tab. 40mg</b> Esomeprazole 40mg	<ul style="list-style-type: none"> <li>* Patients requiring continuous NSAID administration</li> <li>* Treatment of Zollinger-Ellison syndrome</li> <li>* Maintenance therapy after induction of rebleeding prevention for gastric or duodenal ulcers following intravenous treatment</li> </ul>			

## Images

## Products Compositions

## Indications

## Administration · Dosage

## Packing Unit

## ATC Code

### Familan Tab. 20mg

Famotidine 20mg

- \* Gastric ulcer, duodenal ulcer, anastomotic ulcer, upper gastrointestinal bleeding (due to peptic ulcer, acute stress ulcer, hemorrhagic gastritis), reflux esophagitis, Zollinger-Ellison syndrome
- \* Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema) in the following conditions: acute exacerbation of acute gastritis and chronic gastritis

- \* Gastric ulcer, duodenal ulcer, anastomotic ulcer, upper gastrointestinal bleeding (due to peptic ulcer, acute stress ulcer, hemorrhagic gastritis), reflux esophagitis, Zollinger-Ellison syndrome: 20mg twice daily (after breakfast and after dinner or at bedtime). Alternatively, 40mg once daily (at bedtime).
- \* Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema) in acute exacerbations of gastritis (acute or chronic): 10mg twice daily (after breakfast and after dinner or at bedtime). Alternatively, 20mg once daily (at bedtime).

30T/Bottle  
300T/Bottle

A02BA03

### Newxanthine Cap. 150mg

Nizatidine 150mg

- \* Treatment of active gastric and duodenal ulcers and prevention of duodenal ulcer recurrence
- \* Treatment of heartburn symptoms associated with endoscopically diagnosed erosive and ulcerative esophagitis, and gastroesophageal reflux disease (GERD)
- \* Improvement of gastric mucosal lesions (erosion, hemorrhage, erythema, edema) in the following conditions: acute gastritis and acute exacerbation of chronic gastritis

- \* Treatment of active gastric and duodenal ulcers: 300mg once daily at bedtime, or 150mg orally twice daily (morning and evening). Most ulcers heal within 4 weeks, but treatment may be extended up to 8 weeks if necessary.
- \* Prevention of duodenal ulcer recurrence: 150mg orally once daily at bedtime after duodenal ulcer healing
- \* Gastroesophageal reflux disease (GERD): 150mg orally twice daily (morning and evening)
- \* Improvement of gastric mucosal lesions: 150mg orally twice daily (morning and evening)

30C/Bottle  
300C/Bottle

A02BA04

### Newtoga Tab. 10mg

Lafutidine 10mg

- \* Treatment of stomach and duodenal ulcers
- \* Improvement of gastric mucosal lesions in acute gastritis and acute exacerbation of chronic gastritis
- \* Antibiotic combination therapy for peptic ulcer patients infected with Helicobacter pylori
- \* Pre-anesthetic medication
- \* Treatment of reflux esophagitis

- \* Short-term treatment of gastric ulcer: 10mg orally, take twice daily (morning and before bedtime). Most cases are resolved within the prescribed treatment period.
- \* Short-term treatment of duodenal ulcer: 10mg orally, take twice daily (morning and before bedtime). Treatment duration may vary based on patient response.
- \* Improvement of gastric mucosal lesions in acute gastritis and chronic gastritis: 10mg orally once daily, take before bedtime.
- \* Antibiotic combination therapy for peptic ulcer patients infected with Helicobacter pylori: Lafutidine 20mg, Clarithromycin 500mg, and Amoxicillin 1,000mg, take twice daily for 7 to 14 days.
- \* Pre-anesthetic medication: 10mg orally, take twice (once before bedtime on the day prior to surgery and again 2 hours before anesthesia on the day of surgery).
- \* Treatment of reflux esophagitis: 10mg orally, take twice daily.

30T/Bottle  
300T/Bottle

A02BA08

### Mucoran Tab.

Rebamipide 100mg







- \* Gastric Ulcer
- \* Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema) in the following conditions: acute gastritis and acute exacerbation of chronic gastritis

- \* Take 100mg per dose, three times daily.
- \* For gastric ulcer, take in the morning, evening, and before bedtime.

30T/Bottle  
500T/Bottle

A02BX14









Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>NP Ren Tab.</b> Artemisia herb 95% ethanol soft extract (20→1) 60mg	<ul style="list-style-type: none"> <li>* Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema) in the following conditions: acute gastritis and chronic gastritis</li> <li>* Prevention of gastritis induced by the use of nonsteroidal anti-inflammatory drugs (NSAIDs)</li> </ul>	Take 60mg per dose, three times daily.	30T/Bottle 500T/Bottle	A02X
	<b>NP Ren S Tab.</b> Artemisia herb 95% ethanol soft extract(20→1) 90mg	Improvement of gastric mucosal lesions (erosion, bleeding, redness, edema) in the following conditions: acute gastritis and chronic gastritis	Take 90mg per dose, twice daily after meals.	30T/Bottle 300T/Bottle	A02X
	<b>Tiparone Tab.</b> Tiropamide HCl 100mg	<ul style="list-style-type: none"> <li>* Acute spasmodic pain associated with:                             <ol style="list-style-type: none"> <li>1. Hepatobiliary colic</li> <li>2. Abdominal colic of various etiologies</li> <li>3. Renal and ureteral colic</li> </ol> </li> <li>* Abdominal cramps and pain in the following conditions:                             <ol style="list-style-type: none"> <li>1. Gastrointestinal motility disorders</li> <li>2. Cholelithiasis</li> <li>3. Cholecystitis</li> <li>4. Postoperative adhesions</li> </ol> </li> </ul>	Take 100mg per dose, 2 to 3 times daily.	30T/Bottle 500T/Bottle	A03AC05
	<b>Levoran Tab.</b> Levosulpiride 25mg	Relief of the following symptoms associated with functional dyspepsia: bloating, epigastric discomfort, heartburn, belching, nausea, and vomiting	Take 25mg per dose, three times daily before meals.	30T/Bottle 500T/Bottle	A03FA
	<b>Mosagatin Tab.</b> Mosapride citrate hydrate 5mg	Digestive symptoms associated with functional dyspepsia, including heartburn, nausea, and vomiting	15mg per day, divided into three doses and take orally before or after meals.	30T/Bottle 300T/Bottle	A03FA09
<b>NEW</b>					
	<b>Acitoba Cap.</b> 300mg Lactobacillus acidophilus 300mcg	Normalization of bowel movements, constipation, loose stools, abdominal bloating, and abnormal intestinal fermentation	<ul style="list-style-type: none"> <li>* Adults and children 12 years and older: 300mg (1 x 10<sup>8</sup> CFU) 1-3 times daily.</li> <li>* Children 8 to 11 years old: 300mg (1 x 10<sup>8</sup> CFU) 1-2 times daily.</li> </ul>	300C/Bottle	A07FA01



# Urinary System


G

04

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Newfecin Tab. 5mg</b> Solifenacin succinate 5mg	Treatment of overactive bladder symptoms such as urge incontinence and frequent urination	* 5mg once daily.  * If necessary, increase to a maximum of 10mg once daily.	100T (10T/PTP*10) /Pack	G04BD08
	<b>Talosin SR Tab. 0.2mg</b> Tamsulosin HCl 0.2mg	Urinary symptoms associated with benign prostatic hyperplasia (BPH)	Take 1 tablet once daily.	30T/Bottle 100T/Bottle	G04CA02
	<b>Talosin SR Tab. 0.4mg</b> Tamsulosin HCl 0.4mg				
	<b>Tetrin Tab. 2mg</b> Terazosin HCl 2.347mg (as terazosin 2mg)	* Lower urinary tract symptoms associated with benign prostatic hyperplasia (BPH)  * Mild to moderate hypertension	* First dose for dysuria associated with benign prostatic hyperplasia (BPH): 1. Initial dose: 1mg orally before bedtime. 2. Maintenance dose: 5 to 10mg orally once daily.  * High blood pressure: 1. Initial dose: 1mg orally before bedtime. 2. Maintenance dose: 2 to 10mg once daily, take in the morning or evening.	30T/Bottle 100T/Bottle	G04CA03
	<b>Enpiscar Tab.</b> Finasteride 5mg	* Improvement of symptoms of benign prostatic hyperplasia (BPH): 1. Reduction of prostate enlargement and improvement of urine flow 2. Lowering the risk of acute urinary retention 3. Reducing the need for surgical procedures, including transurethral resection of the prostate (TURP) and prostatectomy	* Take 1 tablet once daily, with or without food.  * Even if symptoms improve during treatment, a minimum duration of 6 months is required.	30T (10T/PTP*3) /Pack	G04CB01
	<b>NP-dart Tab.</b> Dutasteride 0.5mg	* Treatment of benign prostatic hyperplasia (BPH): 1. Improvement of symptoms associated with BPH 2. Reduction in the risk of acute urinary retention 3. Decrease in the need for surgical interventions, such as transurethral resection of the prostate (TURP) or prostatectomy 4. Treatment of androgenetic alopecia in adult males (ages 18 to 50): Promotes hair regrowth and slows hair loss progression	* Take 1 tablet once daily.  * Swallow whole without chewing or splitting, after meals.	30T (10T/PTP*3) /Pack	G04CB02

# Adrenal System

H 02


Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Hytisone Tab.</b> Hydrocortisone 10mg	<ul style="list-style-type: none"><li>* Endocrine disorders:<ol style="list-style-type: none"><li>1. Primary and secondary chronic adrenocortical insufficiency</li><li>2. Acute adrenocortical insufficiency</li><li>3. Subacute thyroiditis</li><li>4. Malignant exophthalmos associated with thyroid disease</li><li>5. Adrenocorticotrophic hormone (ACTH) deficiency</li><li>6. Idiopathic hypoglycemia</li></ol></li><li>* Rheumatoid disorders:<ul style="list-style-type: none"><li>- As adjuvant therapy for short-term use to prevent acute progression or exacerbation in the following conditions: chronic rheumatoid arthritis rheumatoid arthritis (including rheumatoid heart disease)</li></ul></li><li>* Collagen diseases:<ol style="list-style-type: none"><li>1. Systemic lupus erythematosus (SLE)</li><li>2. Systemic vasculitis (including aortic syndrome, periarteritis nodosa, and polyarteritis)</li><li>3. Polymyositis (including dermatomyositis)</li><li>4. Scleroderma</li></ol></li></ul>	Take 10 to 120mg per day, divided into 1 to 4 doses.	30T/Bottle 500T/Bottle	H02AB09

Adrenal System

Antiviral Agents

# Antiviral Agents

J 05

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Pamdiac Tab.</b> Famciclovir 250mg	<ul style="list-style-type: none"><li>* Herpes zoster (shingles) infection</li><li>* Treatment of genital herpes infection and suppression of recurrent genital herpes outbreaks</li></ul>	<ul style="list-style-type: none"><li>* Adults:<ol style="list-style-type: none"><li>1. Treatment of herpes zoster infection: 250mg orally 3 times a day for 7 days</li><li>2. Treatment of primary genital herpes infection: 250mg orally 3 times a day for 5 days</li><li>3. Treatment of acute recurrent genital herpes infection: 125mg orally twice a day for 5 days</li><li>4. Suppression of recurrent genital herpes: 250mg orally twice a day</li></ol></li></ul>	21T/Bottle	J05AB09

Product List

## Images

Products  
Compositions

## Indications

## Administration · Dosage

## Packing Unit

ATC  
Code**Amoctam Tab.  
500mg**Amoxicillin 250mg /  
Pivoxil sulbactam 250mg

\* Upper respiratory tract infections: otitis media, sinusitis, tonsillitis, pharyngitis, laryngitis

\* Lower respiratory tract infections: bronchitis, pneumonia, upper and lower urinary tract infections, skin and wound infections, gynecological infections, abdominal infections

Take 1 to 2 tablets per dose, orally administered three times daily.

30T/Bottle  
300T/Bottle

J01CR02

**Amokcle Tab.  
375mg**Amoxicillin hydrate 250mg /  
Potassium clavulanate  
diluted 212.73mg  
(as potassium clavulanate  
125mg)

Acute, chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, urethritis, pyelonephritis, pelvic infection, gonorrhea, boils and abscesses, cellulitis, wound infection, osteomyelitis, dental infection

\* Adults and children aged 12 years or older, or weighing over 40kg: 250mg/125mg orally, three times daily every 8 hours

80T  
(8T/PTP\*10)  
/Pack**Amokcle Tab.  
625mg**Amoxicillin hydrate  
500mg / Potassium  
clavulanate diluted  
212.73mg(as potassium  
clavulanate125mg)

\* For severe infections or respiratory infections, the dose may be increased to 500mg/125mg.

40T  
(4T/PTP\*10)  
/Pack

J01CR02

**Amokcle Syrup  
(7:1)**Amoxicillin hydrate  
36.3143g /  
Potassium clavulanate  
diluted 5.1878g

\* For mild to moderate infections: 25mg/3.6mg per kg of body weight daily, divided into two doses

\* For severe infections: 45mg/6.4mg per kg of body weight daily, divided into two doses

50mL/Bottle

**Safdin Cap.  
500mg**

Cephadrine hydrate 500mg

Tonsillitis, pharyngitis, lobar pneumonia, bronchitis, cystitis, pyelonephritis, urethritis, prostatitis, abscess, cellulitis, boils, impetigo, otitis media

\* Skin and soft tissue infections, respiratory infections: 250 to 500mg orally four times daily, or 500mg to 1g orally twice daily

\* Urinary tract infection: 1 capsule (strength unspecified) orally twice daily

30C/Bottle  
500C/Bottle

J01DB09

**Cefler Cap.**

Cefaclor 250mg






Otitis media, pneumonia, pharyngitis, tonsillitis, bronchitis, pyelonephritis, cystitis, gonococcal urethritis, swelling, carbuncle, folliculitis, cellulitis, infectious atheroma, subcutaneous abscess, whitlow, wound infection

\* Take 1 capsule every 8 hours.

\* For severe infections, take 2 capsules per dose.

30C/Bottle  
500C/Bottle



J01DC04



Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Roxitirocin Tab.</b> Roxithromycin 150mg	Respiratory infections such as pharyngitis, acute bronchitis, tonsillitis, bacterial pneumonia, mycoplasma pneumonia, genital infections and sexually transmitted diseases excluding infections caused by gonorrhea, otitis media, sinusitis, folliculitis, boil, edema, carbuncle, erysipelas, cellulitis, lymphangitis (lymphadenitis), whitlow, paronychia suppurative, subcutaneous abscess, hidradenitis, acne scum, infectious atheroma, for the purpose of preventing infection in case of contact with a patient with meningococcal meningitis, pericoronitis, periodontitis	<ul style="list-style-type: none"> <li>* Take 150mg (titer) twice daily before meals in the morning and evening.</li> <li>* For pneumonia, take 300mg (titer) once daily before breakfast.</li> </ul>	30T/Bottle 500T/Bottle	J01FA06
	<b>Clamacin Tab.</b> Clarithromycin 250mg	<ul style="list-style-type: none"> <li>* Lower respiratory tract infections: bronchitis, pneumonia, etc.</li> <li>* Upper respiratory tract infections: pharyngitis, sinusitis, etc.</li> <li>* Skin and skin tissue infections</li> </ul>	<ul style="list-style-type: none"> <li>* Adults:               <ol style="list-style-type: none"> <li>1. Standard dose: 250mg twice daily</li> <li>2. Severe infections: 500mg twice daily</li> </ol> </li> <li>* Mycobacterial infections: 500mg twice daily (in combination with other antimycobacterial agents)</li> </ul>	30T/Bottle 100T/Bottle	J01FA09
	<b>Clamacin Tab.</b> 500mg Clarithromycin 500mg	<ul style="list-style-type: none"> <li>* Mycobacterium avium, Mycobacterium infection caused by Mycobacterium intracellular</li> <li>* Eradication of Helicobacter pylori in patients with duodenal ulcer</li> </ul>	<ul style="list-style-type: none"> <li>* Eradication of Helicobacter pylori in patients with duodenal ulcer: 500mg three times daily with omeprazole</li> <li>* Children under 12 years of age: Take clarithromycin oral suspension</li> </ul>	30T/Bottle 200T/Bottle	
	<b>Clamacin Dry Syrup</b> 125mg/5mL Clarithromycin coated gr. 84.45mg/g (as clarithromycin 35.47mg/g)	<ul style="list-style-type: none"> <li>* Lower respiratory tract infections: bronchitis, pneumonia, etc.</li> <li>* Upper respiratory tract infections: pharyngitis, tonsillitis, sinusitis, etc.</li> <li>* Acute otitis media</li> </ul>	<ul style="list-style-type: none"> <li>* Take 7.5mg/kg of body weight every 12 hours (15mg/kg/day), regardless of meals. The usual duration is 5 to 10 days.</li> <li>* Mycobacterial infections in children: Take 7.5 to 15mg/kg of body weight every 12 hours (15 to 30mg/kg/day).</li> </ul>	100mL/Bottle	J01FA09
	<b>Clamacin Dry Syrup</b> 250mg/5mL Clarithromycin coated gr. 153.54mg/g (as clarithromycin 64.49mg/g)	<ul style="list-style-type: none"> <li>* Skin and skin tissue infections</li> <li>* Mycobacterial infection caused by Mycobacterium avium or Mycobacterium intracellular</li> </ul>	<ul style="list-style-type: none"> <li>* Do not exceed a maximum dose of 500mg per dose, take twice daily.</li> <li>* Treatment must be combined with other antimycobacterial agents.</li> </ul>	70mL/Bottle	

# Antibiotics

J

01

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Eprosin Tab.</b> Ciprofloxacin HCl 291mg (as ciprofloxacin 250mg)	Respiratory infections excluding acute bronchitis, community-acquired pneumonia, ear, nose, and throat infections (excludes preoperative use for pharyngitis, tonsillitis, laryngitis, otitis externa, chronic cholesteatomatous otitis media and chronic otitis media with metastases to bone), bacterial prostatitis, acute simple cystitis, acute pyelonephritis, complex urinary tract infection, gonococcal cervicitis and gonococcal urethritis, gastrointestinal infection, biliary tract infection, skin and soft tissue infection, wound, bone and joint infection, obstetric gynecological infection (except vaginal infection), peritonitis, lacrimal cystitis, pharyngoma (stye), tarsadenitis	<ul style="list-style-type: none"> <li>* Standard dosage: 250 to 500mg orally twice daily, take between meals.</li> <li>* For severe complex infections: 750mg twice daily</li> <li>* Acute infections: Usually 5 to 10 days. Continue treatment for at least 3 days after symptoms resolve.</li> </ul>	30T/Bottle 500T/Bottle	J01MA02
	<b>Levicin Tab.</b> Levofloxacin hydrate 100mg	Complex skin and soft tissue infections, chronic bronchitis, diffuse panbronchiolitis, secondary infection of chronic respiratory disease, community-acquired pneumonia, bronchiectasis (when infected), acute pyelonephritis, cystitis, prostatitis, gonococcal urethritis, nongonococcal urethritis, epididymitis, bacterial dysentery, enteritis, adnexitis, intrauterine infection, bartholinitis, blepharitis, sty, dacryocystitis, tarsadenitis, corneal ulcer, otitis media (excluding preoperative use for chronic cholesteatomatous otitis media and chronic otitis media metastasized to bone), sinusitis, periodontitis, maxillitis, cholecystitis, cholangitis	<ul style="list-style-type: none"> <li>* Standard dosage: 100mg orally 2 to 3 times daily</li> <li>* For severe cases or patients with insufficient response: 200mg orally 3 times daily</li> </ul>	30T/Bottle 200T/Bottle	J01MA12

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Fuzole Cap.</b> Fluconazole 50mg	<ul style="list-style-type: none"><li>* Acute or recurrent vulvovaginal candidiasis</li><li>* Systemic candidiasis, including disseminated candidiasis and other invasive forms (peritoneal, endocardial, pulmonary, urinary tract, etc.)</li><li>* Cryptococcal meningitis and extrapulmonary cryptococcosis (e.g., pulmonary, cutaneous) in:<ul style="list-style-type: none"><li>1. Immunocompetent patients</li><li>2. Immunocompromised patients (including those with AIDS)</li><li>3. Organ transplant recipients</li><li>4. Patients receiving immunosuppressive therapies</li></ul></li></ul>	<ul style="list-style-type: none"><li>* Vaginal Candidiasis: Take 150mg orally as a single dose.</li><li>* Mucosal Candidiasis:<ul style="list-style-type: none"><li>1. Oropharyngeal Candidiasis: Take 50mg orally once daily.</li><li>2. Chronic Atrophic Oral Candidiasis: Take 50mg orally once daily.</li><li>3. Other Mucosal Candidiasis: Take 50mg orally once daily. For refractory mucosal candidiasis, increase to 100mg orally once daily.</li></ul></li><li>* Systemic Candidiasis (including candidemia, disseminated candidiasis, and other invasive Candida infections):<ul style="list-style-type: none"><li>1. On the first day, take 400mg orally as a single dose.</li><li>2. From the second day onward, take 200mg orally once daily, with the dose increased to a maximum of 400mg once daily if necessary.</li></ul></li></ul>	30C/Bottle 100C/Bottle	J02AC01
	<b>Icozole Tab.</b> Itraconazole solid dispersions 253mg (as itraconazole 100mg)	<ul style="list-style-type: none"><li>* Candida vaginitis</li><li>* Pityriasis versicolor</li><li>* Dermatophytosis (ringworm):<ul style="list-style-type: none"><li>1. Tinea corporis (ringworm of the body)</li><li>2. Tinea cruris (ringworm of the upper part of the body)</li><li>3. Tinea manuum (ringworm of the hands)</li><li>4. Tinea pedis (athlete's foot or ringworm of the feet)</li></ul></li><li>* Oral candidiasis</li><li>* Fungal keratitis</li><li>* Onychomycosis (hand and toenail fungal infections)</li><li>* Systemic fungal infections, such as:<ul style="list-style-type: none"><li>1. Aspergillosis</li><li>2. Candidiasis</li><li>3. Cryptococcosis (including cryptococcal meningitis)</li><li>4. Paracoccidioidomycosis</li></ul></li></ul>	<ul style="list-style-type: none"><li>* Take immediately after meals.</li><li>* Short-term treatment:<ul style="list-style-type: none"><li>1. Candidal vaginitis: 200mg twice daily for 1 day or 200mg once daily for 3 days</li><li>2. Pityriasis versicolor: 200mg once daily for 7 days</li><li>3. Tinea corporis, tinea cruris, tinea pedis (interdigital): 100mg once daily for 15 days</li><li>4. Tinea manuum, tinea pedis (plantar): 100mg once daily for 30 days or 200mg twice daily for 7 days</li><li>5. Oral candidiasis: 100mg once daily for 15 days</li><li>6. Fungal keratitis: 200mg once daily for 21 days</li></ul></li><li>* Onychomycosis:<ul style="list-style-type: none"><li>1. Pulse therapy: 200mg twice daily for 1 week, followed by 3 weeks without treatment. Repeat for 2 cycles for fingernail infections and up to 3 cycles for toenail infections.</li><li>2. Continuous therapy: 200mg once daily for 3 months</li><li>3. For systemic fungal infections in immunocompromised patients, double the dose.</li></ul></li></ul>	28T/Bottle 100T/Bottle	J02AC02

# Musculoskeletal System





M






03

05

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Epison Tab.</b> Eperisone HCl 50mg	<p>* Painful muscle spasms associated with musculoskeletal disorders: Examples include cervicobrachial syndrome, periarthritis, and low back pain</p> <p>* Spastic paralysis caused by neurological diseases: Conditions include cerebrovascular disorders, spastic myelopathy, cervical spondylosis, postoperative sequelae (including brain and spinal cord tumors), traumatic sequelae (such as spinal cord injury and head trauma), amyotrophic lateral sclerosis (ALS), cerebral palsy, spinal cerebellar degeneration, spinal vascular disorders, subacute spinal optic nerve lesions, and other cerebrospinal diseases</p>	50mg per dose, 3 times a day after meals	30T/Bottle 500T/Bottle	M03BX09
	<b>Epison SR Tab.</b> Eperisone HCl 75mg	Painful muscle spasms associated with musculoskeletal disorders: Cervicobrachial syndrome, periarthritis, and low back pain	Take 1 tablet whole, without crushing, splitting, or chewing, twice a day after meals.	30T/Bottle	M03BX09
	<b>Allobone Plus D Tab.</b> Alendronate sodium 91.37mg (as alendronic acid 70mg) / Cholecalciferol concentrate powder 53.33mg (as vitamin D3 5,600IU)	<p>* Treatment of osteoporosis in postmenopausal women</p> <p>* Treatment of osteoporosis in men</p>	<p>* Take 1 tablet once a week.</p> <p>* Take in the morning with plenty of water, at least 30 minutes before eating, drinking, or taking any other medication.</p>	4T (4T/PTP*1) /Pack	M05BB03
	<b>Serobone Plus Tab.</b> Risedronate sodium 35mg / Cholecalciferol concentrated powder 56mg (as cholecalciferol 5,600 I.U)	<p>* Treatment and prevention of osteoporosis in postmenopausal women</p> <p>* Treatment of osteoporosis in men</p>	Take 1 tablet once a week	4T (4T/PTP*1) /Pack	M05BB07



Images	Products Compositions	Indications	Administration • Dosage	Packing Unit	ATC Code
	<b>Acepoin Tab.</b> Aceclofenac 100mg	Rheumatoid arthritis, ankylosing spondylitis, osteoarthritis (degenerative arthritis) and periartthritis of the scapula, toothache, post-traumatic inflammation, back pain, sciatica, pain due to non-articular rheumatism	Take 100mg per dose, twice a day, every 12 hours.	30T/Bottle 500T/Bottle	M01AB16
	<b>Merokcam Cap.</b> Meloxicam 7.5mg	<ul style="list-style-type: none"> <li>* Short-term symptomatic treatment for acute exacerbation of osteoarthritis (degenerative arthritis) accompanying pain and ataxia</li> <li>* Symptomatic treatment of rheumatoid arthritis</li> <li>* Symptomatic treatment of ankylosing spondylitis</li> </ul>	<ul style="list-style-type: none"> <li>* Take with meals (maximum 15mg per day).</li> <li>* For acute exacerbation of osteoarthritis (degenerative arthritis): Adults take 7.5mg once daily, may increase to 15mg if needed.</li> <li>* For rheumatoid arthritis and ankylosing spondylitis: Adults take 15mg once daily. May be reduced to 7.5mg daily depending on treatment response.</li> </ul>	30C/Bottle 500C/Bottle	M01AC06
	<b>Loxpain Tab.</b> Loxoprofen sodium hydrate 68.1mg (as loxoprofen sodium 60mg)	<ul style="list-style-type: none"> <li>* Anti-inflammatory and analgesic for the following diseases and symptoms: chronic rheumatoid arthritis, osteoarthritis (degenerative arthritis), back pain, periartthritis</li> <li>* Anti-inflammatory and pain relief after surgery, trauma and tooth extraction</li> <li>* Antipyretic and analgesic for the following diseases: acute upper respiratory tract infection (including acute upper respiratory tract infection with acute bronchitis)</li> </ul>	<ul style="list-style-type: none"> <li>* For Indications 1 and 2: 1. Take 60mg orally per dose, 3 times a day. 2. For single therapy, Take 60–120mg orally per dose.</li> <li>* For Indication 3: Take 60mg orally, in principle, 2 times a day, with a maximum daily dose of 180mg.</li> </ul>	30T/Bottle 500T/Bottle	M01AE
	<b>Dexifine Tab.</b> Dexibuprofen 300mg	<ul style="list-style-type: none"> <li>* Chronic polyarthritis, rheumatoid arthritis</li> <li>* Arthrosis</li> <li>* Ankylosing spondylitis</li> <li>* Painful swelling or inflammation after trauma and surgery</li> <li>* Aids in the treatment of infections accompanying inflammation, pain, and fever</li> </ul>	<ul style="list-style-type: none"> <li>* Take 1 tablet orally per dose, 2–4 times a day.</li> <li>* Do not exceed a maximum daily dose of 1,200 mg.</li> </ul>	30T/Bottle 300T/Bottle	M01AE14




Images	Products Compositions	Indications	Administration • Dosage	Packing Unit	ATC Code
	<b>Celecok Cap.</b> 200mg Celecoxib 200mg	<ul style="list-style-type: none"> <li>* Relief of symptoms or signs of osteoarthritis (degenerative arthritis)</li> <li>* Relief of symptoms or signs of rheumatoid arthritis</li> <li>* Relief of symptoms and signs of ankylosing spondylitis</li> <li>* Relief of acute pain in adult (pain after surgery, tooth extraction)</li> <li>* Primary dysmenorrhea</li> </ul>	<ul style="list-style-type: none"> <li>* Osteoarthritis (degenerative arthritis): Take once daily</li> <li>* Rheumatoid arthritis: Take 100mg or 200mg twice daily</li> <li>* Ankylosing spondylitis (AS): Take 200mg once daily, or 100mg twice daily. If no effect is observed after 6 weeks, the dose may be increased up to 400 mg daily. If no effect is observed after 6 weeks of taking 400mg daily, consider alternative treatment options.</li> <li>* Acute pain and primary dysmenorrhea: The initial recommended dose is 400mg. If necessary, an additional 200mg may be administered on the first day. From the second day onwards, if needed, take 200mg twice daily.</li> </ul>	30C/Bottle 100C/Bottle	M01AH01
	<b>Moluming Tab.</b> Morniflumate 350mg	<ul style="list-style-type: none"> <li>* Pain relief, anti-inflammatory for the following diseases and symptoms</li> <li>1. Respiratory diseases (laryngitis, tracheitis, bronchitis)</li> <li>2. Otorhinolaryngology (sinusitis, otitis, tonsillitis, pharyngitis)</li> <li>3. Osteoarthritis</li> </ul>	<ul style="list-style-type: none"> <li>* As the starting dose, take 300mg per dose, twice daily.</li> <li>* Depending on the patient's condition, the dose can be increased to a maximum of 700mg per dose, twice daily.</li> </ul>	30T/Bottle 300T/Bottle	M01AX22
	<b>Mupain SR Tab.</b> Acetaminophen 650mg (IR 325mg, SR 325mg) / Tramadol HCl 75mg		Take 1 tablet per dose, up to 4 tablets per day (Doses should be administered at least 12 hours apart).	30T/Bottle 100T/Bottle	
	<b>Mupain Semi Tab.</b> Acetaminophen 162.5mg / Tramadol HCl 18.75mg	Moderate-severe acute, chronic pain	Take 4 tablets per dose, up to 16 tablets per day (Doses should be administered at least 6 hours apart).	30T/Bottle 300T/Bottle	N02AJ13
	<b>Mupain Semi SR Tab.</b> Acetaminophen 325mg (IR 162.5mg, SR 162.5mg) / Tramadol HCl 37.5mg		Take 2 tablets per dose, up to 8 tablets per day (Doses should be administered at least 12 hours apart).	30T/Bottle 100T/Bottle	

# Nervous System

N

02




06

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Gabaten Cap. 100mg</b> Gabapentin 100mg	<p>* Epilepsy</p> <p>1. Monotherapy (including treatment of patients newly diagnosed with seizures): simple/complex partial seizures with or without secondary systemic symptoms in adolescents or adult aged 13 years or older</p> <p>2. Adjunctive therapy: partial seizures with or without secondary systemic symptoms in children over 3 years of age or adult</p> <p>* Neuropathic pain</p>	<p>* Epilepsy</p> <p>1. Monotherapy and adjunctive therapy for patients 13 years of age or older</p> <ul style="list-style-type: none"> <li>- 1st day: 300mg per day</li> <li>- 2nd day: 600mg per day</li> <li>- From the 3rd day: 900mg per day</li> </ul> <p>2. Adjunctive therapy for children aged 3 to 12 years</p> <ul style="list-style-type: none"> <li>- 1st day: 10mg/kg</li> <li>- 2nd day: 20mg/kg</li> <li>- From the 3rd day: 30mg/kg</li> </ul> <p>* Neuropathic pain</p> <ul style="list-style-type: none"> <li>- 1st day: 300mg per day</li> <li>- 2nd day: 600mg per day</li> <li>- From the 3rd day: 900mg per day</li> </ul>	30C/Bottle 100C/Bottle	<b>N02BF01</b>
	<b>Enpicept Tab. 5mg</b> Donepezil HCl 5mg	Treatment of Alzheimer-type dementia symptoms	<p>* 5mg once daily before bedtime</p> <p>* For 4-6 weeks, take a dose of 5mg.</p>	30T/Bottle	<b>N06DA02</b>
	<b>Enpicept Tab. 10mg</b> Donepezil HCl 10mg		<p>* After evaluating the clinical response during this period, the dose may be increased to 10mg.</p> <p>* For underweight female patients 85 years of age or older: The dose should not exceed 5mg per day</p>		

# Antihistamines

R

06

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Levotizine Tab.</b> Levocetirizine HCl 5mg	Relief of symptoms associated with the following conditions:	Take 1 tablet once daily, with or without food.	30T/Bottle 300T/Bottle	<b>R06AE09</b>
	<b>Bepotatine Tab.</b> Bepotastine besilate 10mg	<p>1. Seasonal and perennial allergic rhinitis (including persistent allergic rhinitis)</p> <p>2. Chronic idiopathic urticaria</p> <p>3. Pruritus associated with dermatitis and eczema (in combination with topical hydrocortisone)</p>			
	<b>Azellas Tab.</b> Azelastine HCl 1mg	Perennial allergic rhinitis, chronic idiopathic urticaria, and pruritus associated with skin conditions (such as eczema and dermatitis)	Take 10mg twice daily.	30T/Bottle 300T/Bottle	<b>R06AX</b>
		Bronchial asthma, allergic rhinitis, chronic urticaria, eczema (including atopic dermatitis), and pruritus associated with skin conditions such as dermatitis.	<p>* Take 1 tablet at a time.</p> <p>* For bronchial asthma, take 2mg orally twice daily, once after breakfast and once before bedtime.</p>	30T/Bottle 300T/Bottle	<b>R06AX19</b>

Nervous System

Antihistamines

Product List

# Respiratory System

R

01

03

05

07

## Images

## Products Compositions

## Indications

## Administration · Dosage

## Packing Unit

## ATC Code



### Hederan Syrup

Hedera fluid extract  
20mg/mL

Improvement of symptoms in chronic inflammatory bronchial disease and relief of acute respiratory tract inflammation accompanied by coughing

- \* Adults and adolescents 15 years and older:  
100mg three times daily
- \* Children 4-14 years:  
100mg twice daily
- \* Children 2-3 years:  
50mg three times daily

500mL/Bottle

R05CA12



### Mucidin Cap. 200mg

Acetylcysteine 200mg

Difficulty in evacuating sputum associated with the following conditions: Acute and chronic bronchitis, bronchial asthma, laryngitis, sinusitis, and cystic fibrosis

- \* Acute Disease
  1. Adults: 200mg three times daily
  2. Children (6-14 years): 200mg twice daily
- \* Chronic Disease
  1. Adults: 200mg twice daily
  2. Children (6-14 years):  
100mg three times daily
- \* Cystic Fibrosis
  - Children over 6 years:  
200mg three times daily

30C/Bottle  
500C/Bottle

R05CB01



### Esten Cap. 300mg

Erdosteine 300mg

Mucolysis and expectoration in acute and chronic respiratory diseases

Take 300mg, 2 to 3 times daily.

30C/Bottle  
500C/Bottle

R05CB15



### Lpizine Tab.

Levodropropizine 60mg

Cough in the following diseases: acute and chronic bronchitis

Adults: Take 60mg three times daily, at least 6 hours apart.

30T/Bottle  
300T/Bottle

R05DB27



### Newkamin Syrup

Pelargonium sidoides 11% ethanol extract (1→8-10) / Glycerin mixed solution (8:2) 171.6mg/1mL

Acute bronchitis

- \* Adults and children over 12 years: 9mL, three times daily
- \* Children aged 6 to 12 years: 6mL, three times daily
- \* Children aged 1 to under 6 years: 3mL, three times daily

500mL/Bottle

R07AX



### Mometal Nasal Spray

Mometasone furoate  
0.5mg/mL

- \* Seasonal allergic rhinitis and annual rhinitis
- \* Symptomatic prophylaxis of moderate to severe seasonal allergic rhinitis, nasal polyps, acute rhinosinusitis without severe bacterial infection

- \* For allergic rhinitis and nasal polyps: Spray twice into each nostril once daily.
- \* For acute rhinosinusitis: Spray twice into each nostril twice daily.

18g  
(140 doses)  
/Bottle

R01AD09



### Montest Tab. 10mg

Montelukast 10mg

- \* Prevention and long-term management of asthma
- \* Relief of symptoms of seasonal and perennial allergic rhinitis

- \* For individuals aged 15 years and older: Take 1 tablet once daily.
- \* For asthma patients: Take the tablet in the evening.
- \* For allergic rhinitis patients: Adjust the dosing time based on the patient's condition.
- \* For patients with both asthma and allergic rhinitis: Take 1 tablet in the evening.




28T  
(7T/PTP\*4)  
/Pack

R03DC03

# Anorectic Agents

A

08




Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>PTM Tab.</b> Phentermine HCl 18.75mg (as phentermine 15mg)	* Patients with obesity (BMI > 30 kg/m <sup>2</sup> )  * Overweight patients (BMI > 27 kg/m <sup>2</sup> ) with additional risk factors	Take 1 tablet once or twice daily.	30T/Bottle 100T/Bottle	A08AA01
	<b>Fastin Tab.</b> Phentermine HCl 37.5mg (as phentermine 30mg)		Take 1 tablet once daily.	30T/Bottle 100T/Bottle	
	<b>Phentin Tab.</b> Phendimetrazine tartrate 35mg		Take 1 or 2 tablets 2-3 times daily.	30T/Bottle 300T/Bottle	

Anorectic Agents

# Fat Absorption Inhibitors

A

08

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Zero B Cap. 60mg</b> Orlistat 60mg	Adjunct therapy for weight loss in combination with exercise, behavior modification, and calorie restriction for patients with obesity (BMI > 30 kg/m <sup>2</sup> ) or overweight patients (BMI > 27 kg/m <sup>2</sup> ) with additional risk factors.	Take 120mg three times daily with a fat-containing meal or within one hour after the meal.	30C/Bottle 300C/Bottle	A08AB01
	<b>Zero B Cap. 120mg</b> Orlistat 120mg				
	<b>Zero B Tab. 120mg</b> Orlistat 120mg			90T (10T/PTP*9) /Pack	

Fat Absorption Inhibitors

Product List

# Metabolism Boosters




A

08

16

R





05

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>D-catin Tab.</b> L-carnitine 330mg	Primary and secondary carnitine deficiency: Myocardial metabolic disorders associated with ischemic heart disease, including angina pectoris and acute myocardial infarction	Take 2-3g per day, divided into 2-3 doses.	90T (10T/PTP*9) /Pack	A16AA01
	<b>Daien Cap.</b> Green tea powder 250mg / Orthosiphon powder 150mg	Weight loss supplements for obesity or overweight	Take 1-2 capsules per dose, three times daily.	300C/Bottle	A08AX
	<b>Cafurin-S Tab.</b> Ephedrine HCl 25mg / Caffeine anhydrous 25mg / Acetaminophen 200mg	Relief of common cold symptoms, including sore throat, cough, sputum production, chills, fever, headache, joint pain, and muscle pain.	Take 1 tablet per dose, three times daily.	60T/Bottle	R05X

# Carbohydrate Absorption Inhibitors

V

03

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Luthione Tab.</b> 50mg Glutathione(Reduced) 50mg	Drug addiction	1-3 times daily (50mg: maximum 6 tablets per day; 100mg: maximum 3 tablets per day)	300T/Bottle	V03AB32
	<b>Luthione Tab.</b> 100mg Glutathione(Reduced) 100mg			100T/Bottle	
	<b>Carboglu-X Cap.</b> White kidney bean powder / Vitamin B / Vitamin B2 / Vitamin B6 / Vitamin C / Whey calcium	Indications 1. Vitamin B1: Essential for carbohydrate and energy metabolism 2. Vitamin B2: Essential for energy production in the body 3. Vitamin B6: Essential for protein and amino acid utilization; helps maintain normal blood homocysteine levels 4. Vitamin C: Essential for: 1) Formation and function of connective tissue 2) Iron absorption 3) Protection of cells from free radicals	Take 2 capsules per dose, twice daily.	300C/Bottle	-
<b>NEW</b>					
	<b>Burning Slim Tab.</b> Garcinia cambogia / SNU wonder habanero / Hydroxypropyl methylcellulose (HPMC)	The inhibition of carbohydrate conversion into fat may help reduce body fat.	Take 2 tablets per dose, twice daily.	180T/Bottle	-








# Weight Loss Medications

A

03

06

08

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Tambutine Tab.</b> 200mg  Trimbutine maleate 200mg	* Esophageal reflux and hiatus hernia, gastroduodenitis, gastric and duodenal ulcers and associated digestive disorders (abdominal pain, indigestion, nausea, vomiting), irritable bowel syndrome  * Pediatric diseases: recurrent vomiting, functional gastrointestinal disorders (constipation, diarrhea), irritability, sleep disturbance	1/2 to 1 tablet at a time, 3 times a day	500T/Bottle	A03AA05
	<b>Mano-L Tab.</b>  Aloe extract 100mg / Magnesium hydroxide 500mg / Ursodeoxycholic acid 10mg	Constipation, relief of the following symptoms due to constipation: loss of appetite, abdominal distension, abnormal intestinal fermentation, hemorrhoids	* Adults and children 15 years and older: Take 1 tablet three times daily.  * Children 11-14 years old: Take 2 tablets twice daily.  * Take with or without food (before meals, between meals, or after meals).	300T/Bottle	A06AB20
	<b>Newcodyl Tab.</b>  Bisacodyl 3mg / Docusate sodium 16.75mg	Relief of the following symptoms due to constipation: loss of appetite, abdominal distension, abnormal intestinal fermentation, hemorrhoids	* Adults and adolescents 15 years and older: Take 1-2 tablets three times daily after meals.  * Children 11-14 years old: Take 2/3 to 1 1/3 tablets three times daily after meals.  * Children 7-10 years old: Take 1/2 to 1 tablet three times daily after meals.	500T/Bottle	A06AB52
	<b>Mermad Tab.</b>  Alginic acid 200mg / Carboxymethyl cellulose sodium 100mg	Adjunct therapy for weight loss through appetite suppression	Take 2 tablets per dose, three times daily.	300T/Bottle	A08AX
	<b>Cambel Tab.</b>  Bangpoongtongsungsan dried extract (4.64→1) 237.5mg	The following symptoms of a patient with a lot of subcutaneous fat in the stomach and constipation: accompanying symptoms of high blood pressure (palpitations, stiff shoulders, flushing), obesity, swelling, constipation	Take 4 tablets per dose, three times daily (before or between meals).	500T/Bottle	A08AX
<b>NEW</b>					
	<b>Cambel Double Tab.</b>  Bangpoongtongsungsan dried extract (4.6→1) 475mg		Take 2 tablets per dose, three times daily (before or between meals).	300T/Bottle	
	<b>Naomi Tab.</b>  Banggihwanggitang dried extract (7.52→1) 244mg	The following symptoms are observed: pale complexion, fatigue, excessive sweating, obesity (characterized by loose muscles and fat accumulation), joint pain, and swelling	4 tablets at a time, 3 times a day (Take before or between meals.)	360T/Bottle	A08AX



# Weight Loss Medications

C

10

N










03

06

07

R

06

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Cholesten Soft Cap.</b> Safflower oil 333.33mg (as linoleic acid 250mg) / Tocopherol acetate 6.67mg / Pyridoxine HCl 0.67mg	Improvement of lipid metabolism due to obesity, diabetic and essential hypercholesterolemia	Take 3 capsules three times daily, 30 minutes before meals.	300C/Bottle	<b>C10B</b>
	<b>Topin Tab. 25mg</b> Topiramate 25mg	Prevention of epilepsy and migraine	* Epilepsy: During the first week, administer 1 tablet in the evening. Gradually increase the dose by 25–50 mg per day every 1–2 weeks, divided into two daily doses.	30T/Bottle 100T/Bottle	<b>N03AX11</b>
	<b>Topit Tab.</b> Topiramate 100mg		* The recommended maintenance dose for monotherapy is 100–200 mg per day.  * Migraine prevention: During the first week, take 1 tablet in the evening. Increase the dose by 1 tablet per day at weekly intervals.		
	<b>Depren Cap. 10mg</b> Fluoxetine HCl 11.2mg (as fluoxetine 10mg)	Depression, bulimia nervosa, obsessive compulsive disorder, premenstrual dysphoric disorder	The daily dose should not exceed 80mg.	30C/Bottle 300C/Bottle	<b>N06AB03</b>
	<b>Depren Cap.</b> Fluoxetine HCl 22.4mg (as fluoxetine 20mg)				
	<b>Narex Tab.</b> Naltrexone HCl 25mg	Used in conjunction with an appropriate management program for the treatment of alcohol dependence and to block the effects of exogenous opioids.	Take 2 tablets once daily.	30T/Bottle 100T/Bottle	<b>N07BB04</b>
	<b>Histan F Tab. 16mg</b> Betahistine HCl 16mg	Vertigo, tinnitus, and fluctuating hearing loss associated with Meniere's disease	* Initial dose: Take 1 tablet three times daily.	30T/Bottle 300T/Bottle	<b>N07CA01</b>
	<b>Histan F Tab. 24mg</b> Betahistine HCl 24mg		* Maintenance Dose: 24–48 mg per day		
	<b>Fine S Tab.</b> Bromelain 45mg / Caffeine anhydrous 20mg / Diphenhydramine HCl 25mg	Relief of symptoms caused by a cold (acute rhinitis), allergic rhinitis or sinusitis	Take 1 tablet three times daily.	300T/Bottle	<b>R06AA52</b>

## Images

## Products Compositions

## Indications

## Administration · Dosage

## Packing Unit

## ATC Code

### NEW



#### NP Vita-B Tab.

Benfotiamine 50mg /  
Pyridoxine hydrochloride  
50mg /  
Folic acid 0.25mg /  
Cyanocobalamin 0.2mg

Supplementation of  
Vitamin B1 and B6  
1. For physical fatigue  
2. During pregnancy and lactation  
3. For reduced physical strength  
during or after illness  
4. The efficacy and effects of the  
vitamins contained in this  
medicine are as follows:  
1) Vitamin B1 (Benfotiamine)  
- Alleviation of the following  
symptoms: neuralgia, muscle pain,  
joint pain (e.g., back pain, stiff  
shoulders), beriberi, and eye  
fatigue  
2) Vitamin B6  
(Pyridoxine Hydrochloride)  
- Alleviation of the following  
symptoms: angular stomatitis  
(cracks at the corners of the  
mouth), cheilitis (inflammation of  
the lips), stomatitis (inflammation  
inside the mouth), glossitis  
(inflammation of the tongue),  
eczema, and dermatitis

For individuals aged 12 years  
and older, including adults:  
Take 1 tablet twice a day or 2  
tablets once a day.

60T/Bottle

A11JA



#### NP Selen-C Tab.

Selenium (from dried yeast)  
66.66mg /  
Ascorbic acid 50mg /  
Zinc oxide 31.11mg /  
Chromium (from yeast)  
12.5mg /  
Manganese sulfate hydrate  
3.07mg

\* Supplementation of Vitamin C  
1. For physical fatigue  
2. During pregnancy and lactation  
3. For reduced physical strength  
during or after illness  
  
\* Supplementation of Zinc  
  
\* Supplementation of Selenium

For individuals aged 12 years  
and older, including adults:  
Take 1 tablet twice a day or 2  
tablets once a day.

60T/Bottle

A11GB



#### Vita B Tab.

Folic acid 250mcg /  
Cyanocobalamin 500mcg /  
Pyridoxine hydrochloride  
50mg

Supplementation of Vitamin B6 is  
recommended in the following  
cases:  
1. During pregnancy and lactation  
2. For relief of the following  
symptoms when physical strength  
is reduced during illness or after  
recovery:  
1) Angular stomatitis (inflammation at  
the corners of the mouth)  
2) Cheilitis (inflammation of the lips)  
3) Stomatitis  
(inflammation of the mouth)  
4) Glossitis  
(inflammation of the tongue)  
5) Eczema  
6) Dermatitis

For children aged 8 years and  
older, and adults: Take 1 tablet  
twice a day.

300T/Bottle

A11JA



#### Aniplus Cap.

Vitamin C / Magnesium  
oxide / Crystalline cellulose /  
L-Tyrosine

1. Vitamin C  
1) Essential for the formation and  
function of connective tissue  
Supports iron absorption  
2) Helps protect cells from free  
radicals  
2. Magnesium  
1) Essential for energy metabolism  
2) Supports the maintenance of  
nerve and muscle function

Take 1 capsule twice daily.

300C/Bottle

-

# Hair Growth Solution

D11

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
--------	--------------------------	-------------	-------------------------	--------------	-------------

	<b>Enpina Tab.</b> Finasteride 1mg	Treatment of androgenetic alopecia in adult men aged 18 to 41	Take orally 1mg once a day, regardless of meals.	30T (10T/PTP*3) /Pack	D11AX10
---	---------------------------------------	---	--	-----------------------------	---------

# Neurological Supplements

N06

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
--------	--------------------------	-------------	-------------------------	--------------	-------------

NEW


	<b>Gitako Tab.</b> 240mg Ginkgo leaf dry extract 240mg	Cognitive decline with symptoms of decreased concentration, memory loss, and dizziness (due to atherosclerosis)	Take 1 tablet once a day, regardless of meals.	300T/Bottle	N06DX02
---	---	---	--	-------------	---------

Image	Product Composition	Indication	Administration · Dosage	Packing Unit	ATC Code
	<b>Multivita Inj.</b> Ascorbic acid 100mg, Biotin 60mcg, Cyanocobalamin 5mcg, Dexpantenol 15mg, Ergocalciferol 5mcg (200IU), Folic acid 400mcg, Niacinamide 40mg, Pyridoxine HCl 4.86mg, Riboflavin sodium phosphate 3.6mg, Thiamine HCl 3.81mg, Tocopherol acetate 10mg, Vitamin A 3,300IU	* Vitamin maintenance therapy for patients receiving intravenous nutrition  * Prevention of vitamin deficiency (excluding vitamin K) in patients with: surgery, severe burns, fractures, severe infections or coma	Dissolve the drug in 5mL of distilled water for injection by gently shaking, then mix with 500 to 1,000mL of intravenous glucose solution, physiological saline solution, or a similar solution within 5 minutes. Administer as an intravenous infusion once daily.	10V/Pack	B05XC
	<b>Vitamin C Inj.</b> 20mL  Ascorbic acid 10g/20mL	Prevention and treatment of vitamin C deficiency in cases of increased vitamin C requirement	50-1,000mg daily, administered via SC, IM, or IV, divided into one or more doses per day.	10V/Pack	A11GA01
	<b>Furmine Inj.</b> 5mg  Fursultiamine HCl 5.46mg/10mL	Prevention and treatment of vitamin B1 deficiency, and supplementation when dietary intake is insufficient due to increased vitamin B1 requirements	Usually 5-100mg daily for adults, administered via SC, IM, or IV.	10A/Pack	A11DA01
	<b>Hexavita Inj.</b>  Pyridoxine HCl 50mg/2mL	Prevention and treatment of vitamin B6 deficiency, and supplementation when dietary intake is insufficient due to increased vitamin B6 requirements	For adults, when oral administration is not feasible, administer 10-100mg of pyridoxine hydrochloride daily, divided into 1-2 doses, via IM, SC, or IV.	50A/Pack	A11HA02
	<b>B-colamin Inj.</b>  Hydroxocobalamin 5mg/2mL	Acute neuritis, polyneuritis, sciatica, rheumatoid neuropathy, cervicobrachial syndrome, myalgia, scapular arthritis, cancer-related neuropathic pain, injury-induced neuralgia, postherpetic neuralgia, and migraine	* Adults: Administer 5,000mcg of hydroxocobalamin once daily via IM or IV.  * Adjust to a lower maintenance dose based on clinical response.	50A/Pack	B03BA03
	<b>Vita-D Inj.</b>  Cholecalciferol 5mg/1mL (200,000IU)	* Prevention and treatment of vitamin D deficiency in adolescents and the elderly  * Treatment of vitamin D deficiency in adults	* Adults: 200,000IU cholecalciferol via IM  * Adolescents: 200,000IU cholecalciferol via IM  * Elderly: 100,000IU cholecalciferol via IM  * The total annual dose of cholecalciferol should not exceed 600,000IU.	10A/Pack	A11CC05

Image	Product Composition	Indication	Administration · Dosage	Packing Unit	ATC Code
	<b>Magnesium Chloride Inj. 10%</b> Magnesium chloride 1g/10mL	<ul style="list-style-type: none"> <li>* Treatment of torsades de pointes</li> <li>* Treatment of acute hypokalemia with hypomagnesemia</li> <li>* Magnesium supplementation for electrolyte imbalance</li> <li>* Magnesium supplementation during parenteral nutrition</li> <li>* Prevention and treatment of preeclampsia</li> </ul>	Treatment of torsades de pointes: Slow IV infusion of 8mmol magnesium (equivalent to 2g magnesium sulfate)	10A/Pack	B05XA11
	<b>Zinc S Inj. 5mL</b> Zinc sulfate hydrate 22mg (as zinc 5mg)/5mL	Zinc supplementation during total parenteral nutrition (TPN)	Follow the doctor's prescription, but monitor plasma trace element levels and use them as a guideline for dosage adjustment.	10V/Pack	B05XA18
	<b>Zinc S Inj. 10mL</b> Zinc sulfate hydrate 44mg (as zinc 10mg)/10mL			10V/Pack	
	<b>Selene Inj.</b> Sodium selenite pentahydrate 1.66mg/10mL	Selenium supplementation for patients with selenium deficiency who cannot obtain adequate intake through diet	<ul style="list-style-type: none"> <li>* Administer 100mcg selenium daily via IM or IV.</li> <li>* If necessary, the dose may be increased up to 200mcg daily.</li> <li>* Monitor serum selenium levels to assess treatment adequacy.</li> <li>* When administered in addition to standard total parenteral nutrition (TPN), the daily dose must be limited to 100mcg of selenium.</li> </ul>	10V/Pack	B05XA20
	<b>Multitrace Inj.</b> Zinc sulfate hydrate 43.9mg / Cupric sulfate 15.7mg / Manganese sulfate 3.08mg / Chromic chloride 205mcg / Selenious acid 327mcg/10mL	Supplementation of trace elements (zinc, copper, manganese, selenium, chromium) during parenteral nutrition by central venous access	Follow the doctor's prescription, but monitor plasma trace element levels and use them as a guideline for dosage adjustment.	10V/Pack	B05XA31

# Antioxidants





Injections

A

16

V

03

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Cindella Inj.</b> Thioctic acid 25mg/5mL	Supplementation in cases of increased thioctic acid demand (e.g., during intense physical activity), Leigh syndrome (subacute necrotizing encephalomyelopathy), ototoxicity (due to streptomycin and kanamycin), and noise-induced (occupational) hearing loss	For adults, administer 10-25mg once daily via IV.	10A/Pack	<b>A16</b> AX01
	<b>Luthione Inj. 600mg</b> Glutathione (reduced) 600mg	Prevention of neurological diseases by cisplatin or similar chemotherapy	In severe cases, administer 600-1,200 mg daily via IM or IV infusion.	10V/Pack	<b>V03</b> AB32
	<b>Luthione Inj. 1200mg</b> Glutathione (reduced) 1,200mg				
	<b>Luthione Inj. 600mg with solvents</b> Glutathione (reduced) 600mg / Water for injection 4mL				

# Circulatory System



Injections

A

16

N

06

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Mesocartin Inj.</b> L-carnitine 2g/10mL	<ul style="list-style-type: none"><li>* Primary and secondary carnitine deficiency</li><li>* Myocardial metabolic disorders due to ischemic heart disease: angina pectoris and acute myocardial infarction</li><li>* Treatment of carnitine deficiency in patients with end-stage renal disease undergoing hemodialysis</li></ul>	Adults: Administer 2-3g of L-carnitine daily, divided into 2-3 doses, via IM or IV.	10A/Pack	<b>A16</b> AA01
	<b>Gitako Inj.</b> Ginkgo leaf extract 17.5mg/5mL	Cerebral circulatory disorders and cerebral dystrophy (cerebrovascular insufficiency), including brain dysfunction characterized by: Decreased cognitive function and alertness, dizziness, tinnitus	Administer 5-10mL daily via IM or IV, or 15mL IV every 2 days.	25A/Pack	<b>N06</b> DX02

Antioxidants

Inj.

Circulatory System

Inj.

Product List

2025

# Spasmolytic

Injections

A

12

Images

Products  
Compositions

Indications

Administration · Dosage

Packing Unit

ATC  
Code



**Mine-M Inj.**

Magnesium sulfate 0.5g/5mL

Convulsions in eclampsia, electrolyte replenishment (for hypomagnesemia), and uterine rigidity (oxytocic effect)

\* 1-5g via IM or IV

\* For electrolyte replenishment: Administer 2-4g daily, divided into several doses, via IM or slow IV infusion.

25A/Pack

A12CC02

# Protein Amino Acid

Injections

G

02

Images

Products  
Compositions

Indications

Administration · Dosage

Packing Unit

ATC  
Code



**Placein M Inj.**

Placenta extract 2mL/2mL

Management of menopausal symptoms

Administer 2mL subcutaneously, three times per week for 2 weeks (total of 6 doses).

10A/Pack

G02CX

# Anti-Malignant-Tumor Agents

Injections

L

03

Images

Products  
Compositions

Indications

Administration · Dosage

Packing Unit

ATC  
Code



**Thymosin Inj.**

Thymosin alpha 1 1.6mg






Adjuvanted influenza vaccines for elderly patients with immunosenescence

Administer 900µg/m<sup>2</sup> (1 vial) with the provided diluent and administer via SC or IM, twice weekly for 4 weeks, starting from the first week of vaccination.

2V  
+ 2A (Solvents)  
/Pack




L03AX





Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Liporase Inj.</b> Hyaluronidase 1,500IU	<ul style="list-style-type: none"> <li>* Enhancement of drug penetration when administered via SC or IM.</li> <li>* Local anesthetic effect when used for subcutaneous injection.</li> <li>* Promotion of reabsorption of excess tissue fluids and blood.</li> </ul>	<p>This medication, 1,500IU, should be mixed with sterile water for injection or normal saline before use.</p> <ol style="list-style-type: none"> <li>1. For subcutaneous infusion (large volume subcutaneous administration)</li> <li>2. For intramuscular injection</li> <li>3. As a local anesthetic</li> <li>4. For extravasation</li> <li>5. Hematoma</li> </ol>	10V/Pack	B06AA03
	<b>New-DN Inj.</b> Polydeoxyribonucleotide sodium 5.625mg/3mL	Management and healing of skin graft donor sites	Administer 1 vial daily by IM or SC.	10V/Pack	D03AX
	<b>Newpien Inj.</b> Sodium polynucleotide 20mg / Lidocaine HCL monohydrate 3.2mg/1mL	To reduce mechanical friction in joints through physical restoration by intra-articular injection	Administer up to 5 times within 6 months (once weekly)	2mL * 5syr /Pack	-
	<b>Collapleo</b> Attelo collagen 3%/1mL	Collagen-containing material used to repair or replace defective or damaged ligaments, tendons, muscles, and biological membranes during medical treatments and surgical procedures	Please refer to the instructions for use.	Prefille syringe (1 tube), 22G * 2ea /Pack	-
	<b>Newfopam Inj.</b> Nefopam HCl 20mg/2mL	For the symptomatic treatment of acute pain, especially postoperative pain	<ul style="list-style-type: none"> <li>* 20mg per dose, administered via IM or IV.</li> <li>* If necessary, repeat administration every 6 hours.</li> <li>* Do not exceed 120mg per day.</li> </ul>	10A/Pack	N02BG06

Musculoskeletal System







Injections

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Enpibone Inj.</b> Ibandronate sodium monohydrate 3.375mg (as ibandronate 3mg)/3mL	Treatment of osteoporosis in postmenopausal women	<ul style="list-style-type: none"><li>* 3mg administered intravenously over 15-30 seconds, once every 3 months.</li><li>* Patients should receive supplemental calcium and vitamin D.</li><li>* If a dose is missed on the scheduled day, administer it as soon as possible, then resume the regular 3-month dosing schedule.</li></ul>	Prefilled syringe (1 tube)/Pack	M05BA06
	<b>Hyaliumplus Inj.</b> Sodium hyaluronate 10mg/1mL	Treatment for knee osteoarthritis and shoulder periarthritis	One syringe administered intra-articularly once a week for 3 consecutive weeks into the knee joint or shoulder joint.	Prefilled syringe (3 tubes)/Pack	M09AX01
	<b>Zoledronic Acid Inj.</b> Zoledronic acid	<ul style="list-style-type: none"><li>* Treatment of Osteoporosis</li><li>1. Reduction in the incidence of hip, vertebral, and non-vertebral osteoporotic fractures in postmenopausal women</li><li>2. Reduction in the incidence of new fractures after low-impact hip fractures</li><li>3. Treatment of osteoporosis in men</li><li>* Prevention of osteoporosis in postmenopausal women</li><li>1. Treatment and prevention of glucocorticoid-induced osteoporosis</li><li>2. Treatment of Paget's disease of bone</li></ul>	<ul style="list-style-type: none"><li>* Intravenous infusion through a vented infusion line at a constant rate:</li><li>1. For the treatment of osteoporosis and treatment/prevention of glucocorticoid-induced osteoporosis: Once yearly</li><li>2. For the prevention of osteoporosis in postmenopausal women: Once every 2 years</li><li>3. For the treatment of Paget's disease of bone: Single intravenous infusion over at least 15 minutes</li></ul>	1 bottle (100mL/Bottle) /Pack	M05BA08

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Vari-L Vaccine Inj.</b>  Each vial contains not less than 2,000 PFU of live attenuated varicella-zoster virus (Oka strain)	Prevention of varicella	Reconstitute with the provided diluent and administer 0.5mL as a single subcutaneous injection.	10V + 10V (Diluents)/Pack	J07BK01
	<b>Agkistrodon Antivenom Inj.</b>  Agkistrodon antivenom 6,000IU	Treatment of viper bite	<ul style="list-style-type: none"><li>* Perform a horse serum hypersensitivity test using a 1:10 dilution (reconstitute this medication with the provided diluent, then further dilute it 10-fold with normal saline).</li><li>* Reconstitute this medication with 20mL of the provided diluent (0.6% sodium chloride injection solution).</li><li>* As soon as possible, administer approximately 6,000 units (20mL) intramuscularly or intravenously, avoiding the area near the bite site.</li><li>* Alternatively, dilute with normal saline and administer intravenously.</li><li>* If symptoms persist, an additional dose of 3,000 to 6,000 units (10 to 20mL) may be given after 2 to 3 hours.</li></ul>	5V + 5V (Diluents)/Pack	J06AA03





## Other Injections

## Injections




Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Licorice Inj. 10mL</b> Ammonium glycyrrhizinate 26.5mg / Glycine 200mg / Cysteine hydrochloride monohydrate 10mg/10mL	Adjuvant therapy for urticaria, eczema, allergic dermatitis, and substance use disorders Improvement of liver function in chronic liver disease	Adults: Standard dose: 2 to 20mL intravenously, once daily. For chronic liver disease: 40 to 60mL intravenously or by intravenous infusion, once daily. When increasing the dose, do not exceed the maximum daily dose of 100mL. The dose should be adjusted appropriately according to age and symptoms.	25A/Pack	A05BA
	<b>Licorice Inj. 20mL</b> Ammonium glycyrrhizinate 53mg / Glycine 400mg / Cysteine hydrochloride monohydrate 20mg/20mL				
	<b>ENG Inj.</b> Adenosine triphosphate disodium trihydrate 21.96mg (as adenosine triphosphate disodium 20mg)/2mL	Myasthenia gravis, heart failure, chronic liver disease with impaired liver function, post-concussion syndrome, and accommodative asthenopia	Adults: 1. The usual dose of adenosine triphosphate disodium salt is 5-20mg administered 1-2 times daily via SC, IM or IV. 2. For intravenous administration, mix with 10-20mL of glucose infusion solution and infuse slowly over 3-5 minutes. 3. The dose should be adjusted appropriately according to age and symptoms.	10A/Pack	A16AX
	<b>Same Inj.</b> S-adenosyl-L-methionine sulfate 192mg (as s-adenosyl-L-methionine 100mg)	Depression	As S-adenosyl-L-methionine: 1. 100-200mg per dose, twice daily for 15-20 days, administered via IV or IM. 2. Reconstitute with the provided diluent before use.	5V + 5A (Diluents)/Pack	N06AX
	<b>Newzenil Inj. 2mL</b> Flumazenil 0.2mg/2mL	Reversal of central nervous system sedation induced by benzodiazepines.	* IV, IV infusion  * For reversal of anesthesia: Initial recommended dose is 0.2mg flumazenil, administered via IV over 15 seconds.	10A/Pack	V03AB25
	<b>Newzenil Inj. 5mL</b> Flumazenil 0.5mg/5mL		* For critically ill patients: Initial recommended dose is 0.3mg flumazenil, administered via IV (Refer to the product manual for detailed instructions.).	5A/Pack	

# Other Injections






## Injections


Images	Products Compositions	Indications	Administration · Dosage	Packing Unit	ATC Code
	<b>Dinewdex Inj.</b> Sugammadex sodium 217.6mg/2mL	Reversal of neuromuscular blockade induced by rocuronium or vecuronium	<p>* This medicine should only be administered by or under the supervision of an anesthesiologist.</p> <p>* The recommended dose of this medicine depends on the degree of neuromuscular blockade induced by rocuronium or vecuronium, regardless of the anesthetic regimen used.</p> <p>* The use of appropriate neuromuscular monitoring methods is recommended to evaluate recovery from neuromuscular blockade. In addition, they should be closely monitored for the occurrence of adverse events, including recurrence of neuromuscular blockade.</p> <p>* This medicine is administered as a single dose by bolus injection. A single dose is infused rapidly (less than 10 seconds) through an existing intravenous line.</p>	10V/Pack	V03AB35
	<b>Enpiflu Premix Inj.</b> Peramivir hydrate 349.4mg (as peramivir 300mg)	Treatment of influenza A or B virus infections in adults and children 2 years of age and older (Treatment should be initiated within 48 hours of the onset of initial symptoms of influenza infection.)	<p>* This medication is a premixed formulation.</p> <p>* One bag (300mg/60mL) is used as a single dose for adults, children 2 years of age and older, and patients with renal impairment, only when it can substitute the appropriate dosage.</p> <p>* Do not divide or dilute the contents.</p>	60mL/Bag	J05AH03
	<b>Water For Injection</b> Water for injection 400mL or Water for injection 800mL	Solvent for various injectable preparations	Use an appropriate amount as a solvent.	400mL/bag, 800mL/bag	V07AB
	<b>DHNP Arginine Premix Inj.</b> L-arginine hydrochloride 10g/100mL	<p>For the acute exacerbation of hyperammonemia caused by the following conditions, when oral administration is not possible, to lower blood ammonia levels:</p> <p>Congenital urea cycle disorders:</p> <ol style="list-style-type: none"> <li>1. Carbamyl phosphate synthetase (CPS) deficiency, ornithine transcarbamylase (OTC) deficiency, argininosuccinate synthetase (ASS) deficiency, argininosuccinate lyase (ASL) deficiency</li> <li>2. Lysinuric protein intolerance</li> </ol>	Administer 0.2 to 1g of L-arginine hydrochloride per kg of patient's body weight as a slow intravenous infusion over at least 1 hour.	100mL/bag	B05XB01

# Anesthetics

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit
	<b>Cindella Lidocaine Plus+ Cream</b> Lidocaine 105.6mg/g	Decreased tactile sensitivity of the male genitalia	Apply 0.25 to 0.66g topically.	500g/Jar
	<b>Lidocan Cream</b> Lidocaine 96mg/g	Decreased tactile sensitivity of the male genitalia	Apply 0.25 to 0.66g topically.	500g/Jar
	<b>Pleincell Cream</b> Lidocaine 25mg/g / Prilocaine 25mg/g	Normal intact skin for local analgesia, genital mucous membranes for superficial minor surgery, and as pretreatment for infiltration anesthesia	Apply the cream on genital areas (mucosa and skin) or leg ulcers should be performed only by or under the direct supervision of a healthcare professional.	450g/Jar



# Fillers

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit
	<b>Cindella Collagen Boost</b> PLLA 154mg / CMC 46mg	Skin rejuvenation	Mix 5~8mL of sterile saline or sterile distilled water with the product, mix by hand for 10~15 minutes or use a mixer for 10 minutes, then inject using a 25G cannula (length 38mm or 50mm).	1V/Pack
	<b>Cindella Collagen Boost L</b> PDLLA 170mg / HA 30mg			
	<b>Pleincell</b> Hyaluronic acid 24mg / lidocaine 0.3%/1mL	Skin rejuvenation and facial aesthetics	Please refer to the instructions for use.	Prefilled syringe 30G X 2ea/Pack
				Prefilled syringe 27G X 2ea/Pack
				Prefilled syringe 25G X 2ea/Pack





Images	Products Compositions	Indications	Administration · Dosage	Packing Unit
	<b>Cindella Collagen Glow</b>  PLA / 50 kinds of nutrients	Skin rejuvenation	<ul style="list-style-type: none"><li>* Remove the caps from both the stimulator and booster.</li><li>* Add the booster to the stimulator and mix well.</li><li>* Use an MTS device or similar tool across the entire face.</li><li>* Apply the mixed solution evenly over the whole face.</li></ul>	1V (Booster) + 1V(Stimulator)/Pack



# Health Supplements

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit
	<b>Ascorbic Acid 3g</b> Ascorbic acid 3g	Supplementation of ascorbic acid	Take with water or other suitable beverage.	30 sachets/Pack 100 sachets/Pack
	<b>MSM</b> MSM 2g	It may support joint and cartilage health.	Take one sachet with water, once a day. Take 1 tablet with water, once a day.	60 sachets/Pack 120T/Bottle
	<b>Super Arginine</b> L-arginine 5,000mg / L-carnitine / Pantothenic acid / SNU wonder / 4 types of vitamin B / 9 types of amino acids	Provides L-arginine and others	Take 1 sachet with water, once a day.	30 sachets/Pack
	<b>Glutamine</b> L-glutamine 6,000mg / L-arginine 1,000mg / Zinc / Vitamin A / Lactoferrin concentrate / Milk protein hydrolysate	Provides L-glutamine and others	Take 1 sachet with water, once a day.	30 sachets/Pack
	<b>Immune Alpha</b> Zinc 2.55mg / Sheep thymus powder / MSM / Salmon milt protein extract powder	* Necessary for normal immune function. * Required for normal cell division.	Take 1 capsule with water, twice a day.	60C (6C/PTP*10) /Pack
	<b>Mono Mega 3</b> EPA 560mg / DHA 250mg / DPA 68mg / Vitamin E 2.4mg	* May help improve blood triglyceride levels and blood circulation. * May help improve dry eyes and contribute to eye health.	Take 2 capsules once daily with water.	60C (10C/PTP*6) /Pack
	<b>Ultimate Q10</b> Coenzyme Q10 100mg / PQQ / Bio perine / Organic olive oil	May help with antioxidant effects and reducing high blood pressure.	Take 1 capsule once daily with water.	30C/Bottle
	<b>Homocystrol B-Complex</b> Vitamin B1 75mg / Vitamin B2 75mg / Niacin 75mg / Pantothenic acid 75mg / Vitamin B6 75mg / Biotin 155mcg / Folate (active form) 400mcg / Vitamin B12 300mcg / Choline 50mg / Inositol 37.5mg / Bio perine 2.5mg	Provides B-complex vitamins	Take 1 capsule once daily with water.	60C/Bottle

# Health Supplements

Images	Products Compositions	Indications	Administration · Dosage	Packing Unit
	<b>Ultra Coral Cal Mag Zinc D</b> Calcium (Coral) 300mg / Magnesium 150mg / Zinc 18mg / Vitamin D 15mcg-RE / Wolf berry extract 10mg / Shark cartilage 10mg	Provides calcium, magnesium, zinc, and vitamin D	Take 1 capsule once daily with water.	90C/Bottle
	<b>Neutral Phyto-C</b> Vitamin C (calcium) 1,000mg / Naringin extract / Hesperidin / Rutin / Rosa canina fruit extract	<ul style="list-style-type: none"><li>* Essential for antioxidant action to protect cells from harmful oxygen species.</li><li>* Crucial for the formation and maintenance of connective tissue.</li><li>* Necessary for iron absorption.</li></ul>	Take 1 capsule once daily with water.	90C/Bottle
	<b>Paraotics</b> Probiotics 10,000,000,000CFU	It may promote lactobacillus proliferation, facilitate smooth bowel movements, and support intestinal health.	Take 1 capsule with water, twice a day.	60C/Bottle
	<b>Vitamin D</b> Vitamin D3 mixture 5,000IU	<ul style="list-style-type: none"><li>* It is essential for the absorption and utilization of calcium and phosphorus.</li><li>* It is crucial for bone formation and maintenance.</li><li>* It may help reduce the risk of osteoporosis.</li></ul>	Take 1 capsule with water, twice a day.	30 films/Pack



**Daehan Nupharm**  
Instagram



**Daehan  
Nupharm**

**Headquarter(Factory)**

66, Jeyakgongdan 1-gil, Hyangnam-eup, Hwaseong-si, Gyeonggi-do, Republic of Korea

**Daehan Nupharm Industry-University-Research Center**

20, Changeop-ro 57beon-gil, Sujeong-gu, Seongnam-si, Gyeonggi-do, Republic of Korea

[https:// www.dhnp.co.kr/en/](https://www.dhnp.co.kr/en/)