



DAEHAN
new Pharm



DAEHAN
newPharm

품질관리우수업체
 대한뉴팜(주)

Livestock farming

Antibiotics and Antimicrobials

Antibiotics and Antimicrobials for feed addition

Water-soluble antibiotic, antimicrobials

Liquid and ointment



Antibiotics and Antimicrobials

- **Antibiotics and Antimicrobials for feed addition**
Neoflor powder
Neoflor powder 4%
- **Water-soluble antibiotic, antimicrobials**
Ampicin
CY Tilmipho 200
- **Liquid and ointment**
Neoflor 20% oral sol
Enroxacin 20% Oral





Neoflor powder

(네오플러산)

Medication for Swine Respiratory Disease

- **Composition** (Each 1kg Contains)
Florfenicol..... 20g(potency)
- **Characteristics**
 - It has a very high sensitivity to major causative agent of porcine respiratory disease and shows an excellent antibacterial activity compared to other antibiotics.
 - It is a safe agent that does not cause adverse effects in humans, has a superior efficacy in treating both chloramphenicol-sensitive bacteria and chloramphenicol-resistant bacteria.
 - The results of a comparative study on the major serotypes (type 2, 5) of pleuropneumonia demonstrated a superior therapeutic effect at a lower concentration than the other antibiotics tested.
- **Indications** Prevention and treatment of swine respiratory disease caused by florphenicol-sensitive bacteria For pig feed, uniformly mix 1 ~ 2kg (0.1% ~ 0.2%) per ton and administer for 7 days
- **Caution** Withdrawal period : 3days



Neoflor powder 4%

(네오플러산 4%)

Medication for Swine Respiratory Disease

- **Composition** (Each 1kg Contains)
Florfenicol..... 40g(potency)
- **Characteristics**
 - It has a very high sensitivity to major causative agent of porcine respiratory disease and shows an excellent antibacterial activity compared to other antibiotics.
 - It is a safe agent that does not cause adverse effects in humans, has a superior efficacy in treating both chloramphenicol-sensitive bacteria and chloramphenicol-resistant bacteria.
 - The results of a comparative study on the major serotypes (type 2, 5) of pleuropneumonia demonstrated a superior therapeutic effect at a lower concentration than the other antibiotics tested.
- **Indications** Prevention and treatment of the disease below caused by florphenicol-sensitive actinobacillus.
- Pigs : pleuropneumonia
- **Dosage & Administration** For pig feed, uniformly mix 500g~1kg(0.05~0.1%) per ton and administer for 7 days
- **Caution** Withdrawal period : 3days





Ampicicin

(암피신)

Extensive water-soluble antibiotics

- **Composition** (Each 1kg Contains)
Ampicillin trihydrate 100g
- **Indications** Prevention and treatment of various diseases caused by ampicillin-sensitive gram-positive and gram-negative bacteria
 - Fowl : infectious coryza, enteritis, necrotizing dermatitis, pullorum, salmonellosis, fowl cholera, staphylococcus, colibacillosis
 - Pigs : Salmonellosis, swine erysipelas, pleural pneumonia, bronchitis, cystitis, nephritis, necrotizing enterocolitis, colibacillosis
 - Calves : pneumonia, bronchitis, bacterial diarrhea
 - Dogs, cats : dextrose micrococcus/staphylococcus, Streptococcus dermatitis
- **Dosage & Administration** Dissolve the amount below for administration.
 - Calves : 4 ~ 15g of medication per 100kg bodyweight
 - Pigs, dogs, cats : 0.5 ~ 2g of medication per 10kg bodyweight
 - Fowl : 1g of medication per 10kg bodyweight (allow 180g per 1,000 matured fowl)

Dosage Conditions List

		Broiler Chicken 1,000				
Age of the week		1	2	3	4	5
Dosage(g)		60	120	175	220	225

Layer Chicken 1,000

Age of the week	1	2	3	4	5	6	7
Dosage(g)	25	40	50	60	75	90	100
Age of the week	8	9~10	11~12	13~14	15~17	18~20	More than 20
Dosage(g)	110	130	140	150	160	180	200
- **Caution** Withdrawal period : Cattle, pigs - 5days, Chicken - 2days



CY Tilmipho 200

(CY 틸미포 200)

Medicine - Prevention and treatment

- **Composition** (Each 1kg Contains)
Tilmicosin Phosphate 200g(Potency)
- **Characteristics**
 - treatment of chronic respiratory disease and complex chronic respiratory disease (CRD, CCRD) are excellent,
 - Prevention and treatment of swine dysentery and ileitis are excellent,
 - It has a very high sensitivity and a broad-spectrum antibiotics to Gram-positive and some Gram-negative bacteria., 그 외 a spirochete, large virus and an acid-fast bacterium etc.
 - There is almost no tolerance and low toxic substance
- **Indications**
 - Treatment of bacterial diseases sensitive to Tilmicosin
 - PIG : porcine pleuropneumonia, pasteurella pneumonia, mycoplasma pneumonia
- **Dosage & Administration**
 - PIG : Evenly mix 1~2kg daily (200~400g/ton as Tilmicosin) of medication per 1ton and administer orally for 21 days.
- **Caution** Withdrawal period : 7days





Neoflor 20% oral sol (네오플러 20%액)

Medication for poultry salmonellosis, colibacillosis

- **Composition** (Each 1L Contains)
Florfenicol 200g
- **Characteristics**
 - It exerts a powerful antimicrobial effects to resistant bacteria.
 - Excellent body tissue distribution including lung tissue and has a high bioavailability.
 - It is a safe formulation with no side effects, such as aplastic anemia.
 - It is water-soluble liquid, and will not clog the waterer.
- **Indications**
 - Fowl : treatment of salmonellosis and colibacillosis
 - Pigs : prevention and treatment of porcine pleuropneumonia, pasteurella pneumonia, mycoplasma pneumonia, respiratory disease, salmonellosis, streptococcus
- **Dosage & Administration**
 - Fowl : Dilute 0.5mL of medication with 1L of water and administer for five days.
 - Pigs : Dilute 0.5mL of medication with 1L of water and administer for five days or dilute 0.5mL of medication per 10kg bodyweight for 5 days.
- **Caution** Withdrawal period: Chicken – 5days, Pigs – 16days



Enroxacin 20% Oral (엔로삭신 20%액)

Powerful liquid-type poultry synthetic antibacterial

- **Composition** (Each 1L Contains)
Tilmicosin Phosphate200g
- **Indications**
 - Treatment of bacterial diseases sensitive to enrofloxacin
 - Chicken : E.coli, Mycoplasma gallisepticum, Salmonella typhimurium, Heaemophilus paragallinarum
- **Dosage & Administration**

Animal	Administration	Dosage & Administration
Chicken	Oral administration	Dilute 25mL of medication per 100L of water; have 50mg of enrofloxacin per 1L of daily drinking water and orally administer for 3 days (administer for 5 days for salmonellosis)
- **Caution** Withdrawal period: Chicken – 5days



Anticoccidial drug and an insect repellent

- **Anticoccidial drug**
Coccimuel
Coccimuel-S
- **Fly extirpator (Repellent)**
Minerabon Block





Coccimuel

(콕시멀)

Prevention and treatment for broiler chickens a coccidiosis



- **Composition** (Each 1L Contains)
Sulfaquinoxaline sodium ... 100g Trimethoprim 33.4g
Sodium salicylate..... 100g Menadione sodium bisulfite ... 400mg
- **Characteristics**
 - This is the world's first combined prescription medication for coccidiosis treatment.
 - This "solution" has an excellent solubility suitable for liquid administration.
 - The product has been recognized for its superior efficacy in domestic clinical trials.
 - It contains a prescribed component that effectively treats enterohemorrhage, pain, and fever that appear with coccidiosis and thus can be recovered quickly.
 - This formula is safe even when a dosage as high as 50 times the recommended amount is administered.
 - It has a high reliability. • It has short withdrawal period (7 days).
- **Indications** Prevention and treatment for broiler chickens a coccidiosis
- **Dosage & Administration** Dilute 0.75mL of medication per 1L of water and administer for 3 consecutive days. Give regular water for the next 2~3days. When necessary, dilute 0.75mL of medication per 1L of water and administer for 2~3days.
*Daily dosage for broiler chicken (daily dose per 1,000)

Age of the week	1	7	14	21	28	35	42
Dosage(ml)	50	90	180	250	330	400	450

- **Caution** Withdrawal period : 7days



Coccimuel-S

(콕시멀-에스)

Powerful medication for poultry coccidiosis

- **Composition** (Each 1L Contains)
Diclazuril 5g
- **Characteristics**
 - This is a very effective remedy for coccidiosis.
 - There is almost no tolerance and has a short dose-length.
 - It has an excellent solubility in water, and usually does not cause waterer blockage
 - There are no side effects, and has high safety.
- **Indications** Prevention and treatment of coccidiosis caused by E.acervulina of chickens (broilers)
- **Dosage & Administration** Dilute 0.5mL of medication per 1L of water and administer for 2 consecutive days
- **Caution** Withdrawal period: Chicken - 5days





Minerabon Block (미네라본 부록)

Trace mineral supply and fly larvae repellent supplement

■Composition	(Each 1kg Contains)
	Tetrachlorvinphos..... 10.0g
	Sodium chloride 561g
	Calcium iodate 0.25g
	Ferrous sulfate 6.6g
	Zinc sulfate 0.3g
	Manganese sulfate 0.65g
	Magnesium oxide 6.25g
	Cupric sulfate 0.3g
	Cobaltous sulfate 0.3g
	Sodium selenite 0.1mg

- Characteristics
- This is a functional supplement that added the ingredients to kill fly larvae to the existing mineral supplement. (Summer time mineral supply and fly suppression supplement)
 - Suppressing flies with feces, fundamentally exterminating flies.
 - It is more effective when used in combination with regular insecticide sprays.
 - The product has been permitted by the US FDA and EPA, and it is not harmful to animals.

- Indications
- Relief of fly larvae, supplying salts and inorganic substances to cows, goats, sheep, deer and etc. (iodine, iron, zinc, manganese, magnesium, copper, cobalt, selenium, sodium, calcium).

- Dosage & Administration
- Install in a place suitable for the animals to lick easily and let them freely intake

- Caution
- Withdrawal period : None





feed additive & nutrients

- **Nutrient feed additive**

- Cow-Mix
 - Neovita-Rincal
 - Buffer Milk

- **Water-soluble and liquid nutrients**

- Vita Rich

- **Block Supplements**

- Mineral Block

- **Probiotics, Enzymes, and Yeast**

- Labazyme
 - BioSB-Gold

- **Functional supplementary feeding**



Cow-Mix

(카우믹스)

Health supplements for dairy cows and beef cattle

■Composition	(Each 1kg Contains)
Vitamin A 2,500,000IU
Vitamin D ₃ 500,000IU
Vitamin E 1,000IU
Manganese 4,000mg
Iron 5,600mg
Zinc 1,500mg
Copper 375mg
Iodine 140mg
Cobalt 100mg
Magnesium 3,000mg

- Characteristics**
- Easy ingestion with pellet formulation, and there is no loss.
 - It prevents growth retardation such as rickets and osteomalacia, prevents loss due to breeding failure, and it has effects in increasing the flow rate, milk fat, improving the second-class milk, and preventing ketosis.

■Indications Livestock growth promotion, feed efficiency, improving fattening and milking, enhancing anti-disease power, preventing abnormal milk, paturient palaysis, stress, and osteomalacia.

■Dosage & Administration

Indications	Administration
Facilitate Growth Improved feed efficiency	Calf, Foal(0~3month).....6g Cow in Calf(horse), milk cow(horse)13~23g Beef cattle15~25g
When added to formula feed	administered by mixing in ratio of drug 2 ~ 3g per 1kg of feed
anti-disease enhancement, stress prevention, disease and abnormal milk prevention, paturient palaysis and osteomalacia prevention	administered by mixing in ratio of drug 4 ~ 15g per 1kg of feed (2 times formula feed)





Neovita-Rincal

(네오비타린칼)

Multi vitamin, minerals, calcium phosphate formula

■Composition (Each 1kg Contains)	
Vitamin A.....	1,800,000IU
phosphorus	100g
Sodium	30g
Iron.....	3g
Manganese	60mg
Vitamin D ₃	36,000IU
Calcium	320g
Potassium	300mg
Magnesium	600mg

- Characteristics
- There are powder and pellet type.
 - Palatability and utilization is excellent.
 - Phosphorus, calcium, vitamins and minerals have been prescribed in balance.
 - It prevents rickets, osteomalacia, and growth retardation, loss of milk and reproductive disorders, and solves the problems related to mineral deficiency.

■Indications Supplying vitamin A, vitamin D₃, and minerals, and prevention of the consequent deficiency

■Dosage & Administration Evenly mix 0.4 to 0.6% in cattle feed for administration. (Ratio of drug 4 ~ 6g per 1kg of feed)



Buffer Milk

(버퍼밀크)

Multivitamin, mineral, buffering agent

■Composition (Each 1kg Contains)	
Sodium bicarbonate.....	300g
Vitamin A	250,000IU
Vitamin E	100mg
Manganese	2,700mg
Zinc	1,000mg
Iodine	50mg
Sodium chloride	10g
Magnesium oxide	150g
Vitamin D ₃	50,000 IU
Vitamin B ₁	100mg
Iron	3,000mg
Copper	270mg
Cobalt	5mg

- Characteristics
- The excellent buffering effect sodium bicarbonate and magnesium oxide have been ideally prescribed in combination.
 - Sodium bicarbonate and magnesium oxide prescribed in combination prevents the degradation of blood magnesium level due to the addition of sodium bicarbonate alone.
 - When used in early lactation period, milk fat and flow rate increase is even more enhanced.
 - It maintains the normal pH and the balance of microbial flora.

■Indications

- Cattle: - Improved fat and flow - Urolith prevention
- Prevents acidosis, improves digestibility, promotes growth and appetite
- Improvement of the physical condition by mineral supply

■Dosage & Administration Adult cow (500kg basis) : 150 ~ 400g per capita is administered by mixing the feed.





Vita Rich

(비타리치)

Swine Poultry Health Supplements

■ **Composition** (Each 1kg Contains)

Vitamin A	6,000,000 IU
Vitamin B9	100mg
Vitamin D3	2,000,000 IU
Vitamin B12	2,000µg
Vitamin E	3,000 IU
Vitamin H	10mg
Vitamin C	2,000mg
Nicotinamide	5,000mg
Vitamin K3	1,000mg
Calcium pantothenate	5,000mg
Vitamin B1	3,000mg
DL- Methionine	1,000mg
Vitamin B2	5,000mg
L-lysine	2,000mg
Vitamin B6	1,000mg
Choline bitartrate	30,000mg

- **Characteristics**
- Excellent solubility facilitates oral liquid administration.
 - Poultry and swine health promotion and growth-promoting effects are excellent.
 - It increases disease-fighting ability and enhances productivity.
 - Prevention and treatment of various vitamins and amino acids deficiency are excellent.

- **Indications**
- Chicken : spawn enhancement, growth promotion, feed efficiency, stress and various disease prevention, treatment and prevention of vitamin deficiency
 - Cattle, pigs : Anti-stress, skin diseases, osteomalacia, rickets prevention

- **Dosage & Administration** • Chicken : daily dosage per 1000 (gm)

Age of the week	2weeks	3weeks	4weeks	6weeks	8weeks	More 10weeks
Dosage	5	7~8	10	20	40	50

- Pigs : daily dosage for 10 per capita (gm)

A pig	growing pig	A fully growing pig	A breeding pig
5~10	10~20	20~40	50~100

- Cattle : daily dosage for 10 per capita

Calf	Milk Cow	Beef Cow
20~40	100~200	60~100





Mineral Block

(미네랄 부록)

Trace minerals and salt supplement

■Composition (Each 1kg Contains)			
Iodine	150mg	Manganese	200mg
Sulphur	4,000 mg	Cobalt	100mg
Iron	2,000mg	Zinc	100mg
Nickel	50mg	Copper.....	100mg
Magnesium oxide	3,000mg	Calcium	2,000mg
Selenium	40 µg	Sodium chloride	q.s

- Characteristics**
- The content of various minerals is extremely high compared to other products.
 - Preference is excellent.

- Indications**
- Salts and inorganic substances (iodine, manganese, cobalt, iron, zinc, copper, magnesium, sodium, phosphorus, calcium) are supplied to cattle.

- Dosage & Administration**
- Install in a place suitable for the animals to lick easily and let them freely intake 20-80g daily per capita.



Labazyme

(라바자임)

Mixed live enzymes

■Composition (Each 1kg Contains)	
Lactobacillus acidophilus	Not Less than 2.75x 10 ¹⁰ CFU
Streptococcus faecium	Not Less than 8.25x 10 ¹⁰ CFU
Bacillus subtilis	Not Less than 1.1x 10 ¹⁰ CFU
Protease	Not Less than 2,750 CSU
Amylase.....	Not Less than 5,500 SLU
Cellulase.....	Not Less than 27.5 FPU

- Characteristics**
- Proper intestinal flora of livestock is established to promote growth, feed efficiency, and effectiveness of anti-diarrhea.
 - Live bacteria such as lactic acid bacteria and hay bacillus competitively inhibit colony formation of harmful pathogenic bacteria (salmonella, E. coli, etc.).
 - It contains three kinds of excellent lactic acid bacteria digestive enzymes, thus has an excellent growth-promoting effects and diarrhea prevention.

- Indications**
- For calves, pigs, and fowl,
 - prevention and treatment of diarrhea
 - promote growth, weight gain
 - inhibit growth of intestinal pathogenic bacteria
 - inhibit antibiotic-resistant pathogens
 - remove odor of livestock manure

- Dosage & Administration**
- Administer feed by evenly mixing the following dose.
 - milking cows and beef cattle: 2kg per ton of feed
 - Pigs: 2kg per ton of feed - Chicken: 0.5kg per ton of feed





BioSB-Gold

(바이오에스비-골드)

Highly-enriched complex live yeast, probiotics

- **Composition** (Each 1kg Contains)
- Saccharomyces cerevisiaeNot less than 3.0×10^{11} CFU
 - Bacillus subtilisNot less than 4.0×10^9 CFU
 - Biomass Metabolitesq.s

- **Characteristics**
- This is a yeast culture agent that contains large amount of Bacillus subtilis of the existing yeast culture.
 - It contains general nutrients (protein, fat, vitamins, amino acids, etc.) as well as UGF, which has an excellent growth-promoting effects of livestock.
 - It improves feces condition by inhibiting the production of intestinal ammonia gas in the livestock, decreases odor, and enhances the breeding environment. When added to feed, it increases the palatability.

■ **Indications**

Target animal	By stage of growth	Effects
Fowl	Chick, laying hens, broilers	promote growth, increase egg production, livestock fecal odor removal
Pig	piglets, growing pigs, finishing pigs, sows	promote growth, promote digestion, improve growth rate, remove fecal odor
Cattle	calves, beef cattle, milking cow	growth-promoting, improving milk quality and meat quality, feed efficiency, increase flow, removes odor, diarrhea prevention

■ **Dosage & Administration**

Species	Division	Amount of addition per ton of feed
Dairy	calves	1~2kg
	Dry period	1kg
	Milking period	1kg
Fattening cattle	Calves	1 ~2kg
	Beef cattle	1 kg
	Adult cow	1kg
Pig	Piglets	1~2kg
	Foster, pigs	1~2kg
	Sow, pig husbandry	1kg
Chicken	Broiler	1 kg
	Laying	1~2kg



For Fishery Products

Antibiotic, Antimicrobial
Parasites extirpator/Repellent
Feed additive & Nutrients
Probiotics, Enzymes, and Yeast



Antibiotic, antimicrobial

Aqua Amocillin
Aquaterra






Aqua Amocillin

(아쿠아 아모실린)

Extensive penicillin antibiotics for fish



- Composition** (Each 1kg Contains)
 Amoxicilline trihydrate 200g(Potency)
- Characteristics**
 - is a broad-spectrum antibiotics for Gram-positive and Gram-negative bacteria.
 - It the most cost-effective disease for Pasteurella piscicida and furunculosis prevention

Target animal	Effects	Indication
Yellow tail	Prevention and treatment of amoxicillin-sensitive diseases	Pasteurella piscicida and furunculosis prevention, mortality rate improvement

Target animal	Methods	Dosage and administration
Yellow tail	Oral intake	Daily intake of 100~200g of medication per ton; mix in feed 1~2 times per day and administer for 4~7 consecutive days

- Caution** Withdrawal period: Yellow tail – 7days



Aquater

(아쿠아테라)

This is a broad-spectrum antibiotics prevention and treatment.




- Composition** (Each 1kg/1L Contains)
 Oxytetracycline HCl 240g
- Characteristics**
 - An extensive antibiotic that is very effective for gram-positive and gram-negative bacteria.
 - It is water-soluble, and thus can be taken orally and used for baths.
 - It is the most cost-effective disease prevention (cleaning agent) for fish.

- Indications** Treatment and mortality rate reduction for the following fish diseases caused by oxytetracycline-sensitive bacteria.

Target fish	Indication
Yellow tail/Porgy	Vibrio anguillarum
trout	Haemophilus sp., Aeromonas salmonicida, Aeromonas sp., etc.,
eel	Edwardsiella tarda
carp	Haemophilus sp., Aeromonas sp., Flexibacter columnaris
goldfish	Haemophilus sp. Flexibacter columnaris
halibut	Streptococcus sp.

- Dosage & Administration**
 - Oral administration : evenly mix 210g of the product in feed per ton a day.
 - For baths: dilute 5~25ppm (titer) of oxy tetracycline in water and bath for 30~60 minutes (20~105 g of the product per 1 ton of water).

- Caution** Withdrawal period : yellow tail – 20 days
 eel, Porgy, trout, carp – 30 days
 halibut – 40 days



Parasites extirpator/repellent

Aquaprantel



Aquaprantel

(아쿠아프란텔)

Excellent parasiticide for gill distomias



- **Composition** (Each 1kg Contains)
Praziquantel 500g
- **Characteristics**
- This drug contains a high concentration (50%) of praziquantel, which was identified as highly efficient and low toxic substance in a study that compared the effects of different pesticides.
 - It is especially effective for gill distomias and skin distomias that are fatal to fish.
 - It resolves the stress caused by drug administration.
 - It is also suitable for other drug-resistant parasitic infections.
- **Indications**
- Parasiticide for the parasites(flukes, trichuridae) in aquarium fish (gold fish- gold fish, tropical fish-Astronotus)
 - Gill distomias treatment for Jacopever (Rockfish)
- **Dosage & Administration**
- Aquarium fish
 - Oral administration: Mix 300mg of the product in 100g of feed and administer orally for twice a day for 5 days.
 - For baths: dissolve 40mg of the product into 1L of water and bath for 2 hours. Re-bath after 2 weeks.
 - Jacopever (Rockfish)
 - Oral administration: mix 0.2 ~ 0.4g of the product per kg of fish bodyweight in feed, orally administer 1-3 times every other day
 - For baths: dissolve 0.1 ~ 0.2g of the product into 1L of water and bath the subject fish for 4 minutes.
- **Caution** Withdrawal period : rockfish - 10 days

Liver function improvement and immune enhancement

Aqua Gluton
Aqua Livertonic





Aqua Gluton

(아쿠아글루톤)

Complex liver function improving agent



- **Composition** (Each 1kg Contains)

Glutathion 20g

DL-Methionine 300g
- **Characteristics**

 - Dietary liver dysfunction enhancer for farmed fish.
 - Complex liver function enhancer containing more than 2.0% of glutathione and more than 30% of methionine, the amino acid.
 - It is effective in fat metabolism disorder prevention and treatment, and increases the survival rate, improves disease-fighting ability and growth rate.
- **Indications**

Prevention and treatment of ceroid symptoms caused by deteriorated feed and improving liver function of farmed fish including halibut, yellowtail, carp, and leather carp.
- **Dosage & Administration**

Mix 0.5 ~ 1kg of the product per a ton of feed and administer for more than 7 days. Or add 0.25 ~ 1kg of the product per ton of fish body weight and orally administer for more than 7 days.



Aqua Livertonic

(아쿠아리버토닉)

Liver function enhancer by preventing hepatocellular injury



- **Composition** (Each 1kg Contains)

Glutathion 20g
- **Characteristics**

 - Dietary liver dysfunction enhancer for farmed fish
 - It is effective in fat metabolism disorder prevention and treatment, effective in increasing survival rate, growth rate, and disease-fighting ability
- **Indications**

Liver function improvement, prevention and treatment of ceroid symptoms caused by deteriorated feed for farmed fish including halibut, yellowtail, carp, and leather carp
- **Dosage & Administration**

Mix 0.5 ~ 1kg of the product per a ton of feed and administer for more than 7 days. Or add 0.25 ~ 1kg of the product per a ton of fish bodyweight and orally administer more than 7 days.



Nutrients

Bio-C for Fish



Bio-C for Fish

(수산용바이오-씨)

Stabilized Vitamin-C coated with natural vegetable oil

■ **Composition** (Each 1kg Contains)
Vitamin-C(Ascorbic acid) 200g

■ **Characteristics**

- This will enhance the immune system of fish.
- It increases productivity by enhancing digestibility.
- It reduces the toxicity of organic compounds.
- Prevention and treatment of vitamin C deficiency in fish

■ **Indications** Prevention and treatment of vitamin C deficiency in farmed fish

■ **Dosage & Administration** Administer by evenly mixing the following dose in a ton of feed

fish species	Administration in a ton of feed
shrimp	1.0 ~ 1.5kg
salmon	0 .5 ~ 1.0kg
trout	1.0 ~ 1.5kg





Active yeast · Probiotics · Enzymes

Bio SB-Gold
Labazyme



BioSB-Gold

(바이오에스비-골드)

Active yeast · Probiotics

- **Composition** (Each 1kg Contains)
 - Saccharomyces cerevisiae Not less than 3.0×10^{11} CFU
 - Bacillus subtilis Not less than 4.0×10^9 CFU
 - Biomass Metabolites q,s
- **Characteristics**
 - The growth-promoting effect in fish is excellent.
 - It is effective in improving digestibility and feed efficiency.
 - The yeast extract developed in the United States is used, and the bioavailability is high.
 - Addition of yeast culture improves immune function and disease prevention by its immune-enhancing substances such as beta-glucan and mannanoligosaccharides.
- **Indications** Growth promotion, digestion promotion, and feed efficiency improvement
- **Dosage & Administration** Evenly mix 1 ~ 5kg in a ton of feed and administer.



Labazyme

(라바자임)

Mixed live enzymes

- **Composition** (Each 1kg Contains)
 - Lactobacillus acidophilus Not Less than 2.75×10^{10} CFU
 - Streptococcus faecium Not Less than 8.25×10^{10} CFU
 - Bacillus subtilis Not Less than 1.1×10^{10} CFU
 - Protease Not Less than 2,750 CSU
 - Amylase Not Less than 5,500 SLU
 - Cellulase Not Less than 27.5 FPU
- **Characteristics**
 - Fish growth promotion and feed efficiency improvement are outstanding.
 - It enhances disease prevention and immune function.
 - It prevents reduced feed intake caused by stress.
 - It improves water quality.
- **Indications** Digestion promotion, feed efficiency improvement, and mortality rate reduction
- **Dosage & Administration** Evenly mix 0.5 ~ 2.5kg per a ton of feed and administer

